

THE STACKY CONCENTRATION THEOREM

DHYAN ARANHA, ADEEL A. KHAN, ALEXEI LATYNTSEV, HYEONJUN PARK,
AND CHARANYA RAVI

ABSTRACT. We give a sufficient criterion for the Chow or algebraic bordism groups of an algebraic stack, localized at a set of Chern classes of line bundles, to be concentrated in some closed substack. This is a vast generalization of the torus fixed-point localization theorem in equivariant intersection theory, which is the special case of the stack quotient of a scheme X by an action of a torus T . Taking on the one hand an algebraic stack in place of X , we deduce a generalization of torus localization to algebraic stacks. Taking on the other hand any algebraic group G instead of T , we obtain a localization theorem in G -equivariant intersection theory.

Introduction	2
0.1. Main results	2
0.2. Examples, variants, and generalizations	3
0.3. Proof	5
0.4. Localization formulas	6
0.5. Contents of the paper	7
0.6. Acknowledgments	7
1. Preliminaries on stacks	8
1.1. Stacks	8
1.2. Points	8
1.3. Stabilizers	9
1.4. Stabilizers of group actions	9
1.5. Fixed loci	10
2. Intersection theory on stacks	11
2.1. Motivic Borel–Moore homology	11
2.2. Chow, G -theory, bordism	14
2.3. Localization triangle	15
2.4. Projective bundle formula	16
2.5. Reduction to the subtorus	17
3. Concentration	17
3.1. The master theorem	18
3.2. Generic slices	19
3.3. Proof of Theorem 3.5	20
3.4. Proof of Theorem 3.10	20
3.5. The equivariant master theorem	21
3.6. Base change	22
3.7. Non-quasi-compact stacks	23
4. Torus concentration	25
4.1. The statement	25
4.2. Example: fixed loci	27

4.3. Example: vector bundles and abelian cones	27
4.4. Example: Higgs sheaves on curves	29
References	30

INTRODUCTION

0.1. Main results. The starting point of this paper is the *concentration theorem*¹ in equivariant intersection theory. Given a split torus T over a field k and a k -scheme X with T -action, we denote by $A_*^T(X)$ the T -equivariant Chow group of X with rational coefficients. Then we have (see [EG2, Thm. 1]):

Theorem 0.1. *Let X be a scheme of finite type over a field with an action of a split torus T . Let $i : X^T \hookrightarrow X$ denote the inclusion of the T -fixed locus. Then the push-forward map on T -equivariant Chow groups induces an isomorphism*

$$i_* : A_*^T(X^T)_{\text{loc}} \rightarrow A_*^T(X)_{\text{loc}}$$

where the localization is at the set of first Chern classes of nontrivial 1-dimensional representations of T .

This result forms the core of equivariant localization theory; for instance, it leads immediately to the localization formula of Atiyah and Bott [AB] when X is smooth and to the Bott residue formula [Bot] when X is smooth and proper.

In this paper we prove a new type of concentration theorem for general algebraic stacks. Let $A_*(\mathcal{X})$ denote the cycle groups (with rational coefficients) defined in [Kha1] for any algebraic stack \mathcal{X} of finite type over k .² For the quotient stack $\mathcal{X} = [X/T]$, this agrees with the T -equivariant cycle groups $A_*^T(X)$.³ We claim (see Corollary 3.7):

Theorem A (Stacky concentration). *Let \mathcal{X} be an Artin stack of finite type over a field k with affine stabilizers and $\mathcal{Z} \subseteq \mathcal{X}$ a closed substack. Let Σ be a set of line bundles on \mathcal{X} such that for every geometric point x of $\mathcal{X} \setminus \mathcal{Z}$, there exists $\mathcal{L}(x) \in \Sigma$ whose restriction to $B\text{Aut}_{\mathcal{X}}(x)$ is trivial. Then push-forward along $i : \mathcal{Z} \hookrightarrow \mathcal{X}$ induces an isomorphism*

$$i_* : A_*(\mathcal{Z})[\Sigma^{-1}] \rightarrow A_*(\mathcal{X})[\Sigma^{-1}]$$

where $\mathcal{L} \in \Sigma$ acts via the first Chern class.

¹This is often simply called “the localization theorem”. The less ambiguous terminology is due to R. W. Thomason [Tho2] as far as we know.

²That is, $A_n(\mathcal{X})$ is defined as the motivic Borel–Moore homology group $H_{2n}^{\text{BM}}(\mathcal{X}; \mathbf{Q}^{\text{mot}}(-n))$ of [Kha1], and $A_*(\mathcal{X})$ is the direct sum over $n \in \mathbf{Z}$.

³By [KR1, Cor. 6.5], we have $A_n(\mathcal{X}) \simeq A_{n-\dim(T)}^T(X)$.

From this we derive the following generalization of Theorem 0.1 from schemes to Artin stacks (see Theorem 4.1). Given an Artin stack X over k with T -action, we write $A_*^T(X)$ for the T -equivariant cycle groups of X (with rational coefficients).⁴

Theorem B (Torus concentration). *Let X be an Artin stack of finite type over k with T -action. Let $Z \subseteq X$ be a T -invariant closed substack away from which every point has T -stabilizer properly contained in T .⁵ Then push-forward along $i : Z \hookrightarrow X$ induces an isomorphism*

$$A_*^T(Z)_{\text{loc}} \rightarrow A_*^T(X)_{\text{loc}} \quad (0.2)$$

where the localization is as in Theorem 0.1.

0.2. Examples, variants, and generalizations.

0.2.1. *Deligne–Mumford stacks.* When X is a Deligne–Mumford stack, Theorem B was proven by A. Kresch in the case where the torus T is of rank 1 and the field k is algebraically closed field (see [Kre, Thm. 5.3.5]).

Note that, at least when X is separated, $A_*^T(X) \simeq A_*([X/T])$ agrees with Kresch’s cycle groups [Kre]. Indeed, X admits a coarse moduli space M inheriting the T -action and both cycle groups are canonically isomorphic to $A_*^T(M)$ (recall that we are working with rational coefficients).

0.2.2. *Artin stacks with finite stabilizers.* Suppose X is an Artin stack with finite stabilizers; recall that when k is of positive characteristic, X need not be Deligne–Mumford. For example, this applies to moduli stacks of stable objects in enumerative geometry, e.g. moduli of stable maps in Gromov–Witten theory and moduli of stable sheaves in Donaldson–Thomas theory.

As long as X is separated (or more generally has finite inertia), $A_*^T(X)$ still agrees with Kresch’s cycle groups for the same reason as in the Deligne–Mumford case.

0.2.3. *Global quotient stacks.* Suppose G is a linear algebraic group over k and X is an algebraic space of finite type over k with commuting actions of G and T . Suppose $Z \subseteq X$ is a closed subspace invariant under both actions which contains all points whose T -stabilizer is the whole torus T . Then Theorem B applied to $\mathcal{X} = [X/G]$ yields the isomorphism

$$i_* : A_*^{G \times T}(Z)_{\text{loc}} \rightarrow A_*^{G \times T}(X)_{\text{loc}} \quad (0.3)$$

on $G \times T$ -equivariant Chow groups. This generalizes a concentration theorem proven by A. Minets [Min] under a much more restrictive technical condition.

⁴That is, $A_n^T(X)$ is defined as the relative motivic Borel–Moore homology group $H_{2n}^{\text{BM}}([X/T]_{/BT}; \mathbf{Q}^{\text{mot}}(-n))$ of [Kha1], and $A_*^T(X)$ is the direct sum over $n \in \mathbf{Z}$. For X a scheme or algebraic space, this agrees with the Edidin–Graham equivariant cycle groups again by [KR1, Cor. 6.5].

⁵Informally speaking, the condition on Z means that it contains every T -fixed point. See Subsect. 1.4 for the notion of T -stabilizers (which are not the same as the stabilizers of $[X/T]$ when X is not an algebraic space).

0.2.4. *Locally of finite type stacks.* We will also prove a generalization of Theorem B to *locally* of finite type Artin stacks. In fact, the result is false in this generality with the naïve definition of $A_*^T(X)_{\text{loc}}$. In Subsect. 3.7 we will give another definition of the latter that agrees with the naïve localization in the quasi-compact case, but for which concentration does hold. For example, this applies to the moduli stack of Higgs sheaves on a curve, which admits a canonical scaling action of $T = \mathbf{G}_m$. This was one of the motivating examples⁶ (see Theorem 4.12):

Corollary C. *Let k be a field and C a smooth proper and geometrically connected curve over k . Denote by $\underline{\text{Higgs}}$ the moduli stack of Higgs sheaves on C and by $\mathbf{\Lambda}$ the closed substack parametrizing nilpotent Higgs sheaves. Then the map*

$$A_*^T(\mathbf{\Lambda})_{\text{loc}} \rightarrow A_*^T(\underline{\text{Higgs}})_{\text{loc}}$$

is invertible.

This generalizes a result of A. Minets [Min, Cor. 4.3], who considered the substack of *torsion* (rank zero) Higgs sheaves.

0.2.5. *Localizing K-theory classes.* The following variant of Theorem A can be more useful when there is not a sufficient supply of 1-dimensional representations.

With notation as in Theorem A, let $\Sigma \subseteq K_0(\mathcal{X})$ be a set of K-theory classes such that for every geometric point x of $\mathcal{X} \setminus \mathcal{Z}$, there exists an $\alpha(x) \in \Sigma$ whose restriction to $B\text{Aut}_{\mathcal{X}}(x)$ is trivial. Let $\hat{A}_*(\mathcal{X})$ denote the *product* of the cycle groups $A_n(\mathcal{X})$ over $n \in \mathbf{Z}$. Then we will prove that i_* induces an isomorphism

$$i_* : \hat{A}_*(\mathcal{Z})[\Sigma^{-1}] \rightarrow \hat{A}_*(\mathcal{X})[\Sigma^{-1}] \quad (0.4)$$

where $\mathcal{L} \in \Sigma$ acts via the Chern character. See Theorem 3.10.

0.2.6. *General algebraic group actions.* We can specialize stacky concentration to quotients by arbitrary algebraic group actions $[X/G]$. We get the following generalization of Theorem B (see Corollary 3.15):

Corollary D (Equivariant concentration). *Let G be an algebraic group acting on an Artin stack X of finite type over a field k with affine stabilizers. Let $\Sigma \subseteq K_0(BG)$ be a subset of nonzero elements. Let $Z \subseteq X$ be a G -invariant closed substack containing every point x of X such that no element of Σ is sent to zero by $K_0(BG) \rightarrow K_0(B\text{St}_X^G(x))$. Then*

$$i_* : \hat{A}_*^G(Z)[\text{ch}(\Sigma)^{-1}] \rightarrow \hat{A}_*^G(X)[\text{ch}(\Sigma)^{-1}]$$

is an isomorphism.

⁶We thank Y. Soibelman for suggesting it; see [FSS] for an analogous analysis at the level of virtual motives.

0.2.7. *Oriented Borel–Moore homology theories.* Our methods work for general oriented Borel–Moore type homology theories. This includes usual Borel–Moore homology (over $k = \mathbf{C}$) and ℓ -adic Borel–Moore homology (over a field in which ℓ is invertible).

We also prove concentration for algebraic bordism, namely for the stacky and equivariant (higher) algebraic bordism theories defined in [Kha1, Ex. 2.12] and [KR1, §8]. The analogous statement for Levine–Morel algebraic bordism was proven for smooth and projective varieties over a field of characteristic zero by A. Krishna [Kri], and generalized recently in [KP] to (possibly singular) quasi-projective schemes.

Our methods also apply to G-theory (= algebraic K-theory of coherent sheaves). In this paper, we only consider étale G-theory and Borel-equivariant G-theory (in the sense of [KR1, §7]). However, the case of “genuine” G-theory can be treated by the same method and will be done in detail in [KPR].

0.2.8. *Categorification.* In a sequel [KR2] to this paper, two of the authors proved a categorification of the stacky concentration theorem. For example, consider the stable ∞ -category $\mathbf{DM}(X)_{\mathbf{Q}}$ of (\mathbf{Q} -linear) motivic sheaves on an Artin stack X . In the situation of Theorem A, we have the \star -direct image functor $i_* : \mathbf{DM}(\mathcal{Z})_{\mathbf{Q}} \rightarrow \mathbf{DM}(\mathcal{X})_{\mathbf{Q}}$. Categorified concentration asserts that i_* induces an equivalence of stable ∞ -categories

$$i_* : \mathbf{DM}(\mathcal{Z})_{\mathbf{Q}}[\Sigma^{-1}] \simeq \mathbf{DM}(\mathcal{X})_{\mathbf{Q}}[\Sigma^{-1}] \quad (0.5)$$

where the localization is in an appropriate ∞ -categorical sense.

This result is used to obtain concentration for oriented *cohomology* theories, such as motivic cohomology or Betti and ℓ -adic cohomology, where it is the pullback map $i^* : H^*(\mathcal{X}) \rightarrow H^*(\mathcal{Z})$ that induces an isomorphism after localization.

0.3. **Proof.** Let us briefly comment on the proof of the stacky concentration theorem (Theorem A). The cohomological formalism for intersection theory on stacks of [Kha1] plays an essential role in reducing this vastly general statement to a drastically simpler one.

The cycle groups $A_n(\mathcal{X})$ are realized as the motivic cohomology groups

$$A_n(\mathcal{X}) = H^{-2n}(\mathcal{X}, f^! \mathbf{Q}^{\text{mot}}(-n))$$

where $\mathbf{Q}^{\text{mot}}(-n) \in \mathbf{DM}(\text{Spec}(k))$ is the Tate twist in Voevodsky motives, $f : \mathcal{X} \rightarrow \text{Spec}(k)$ is the projection, and $f^! : \mathbf{DM}(\text{Spec}(k))_{\mathbf{Q}} \rightarrow \mathbf{DM}(\mathcal{X})_{\mathbf{Q}}$ is the compactly supported inverse image functor on stable ∞ -categories of (\mathbf{Q} -linear) motivic sheaves. One can think of the right-hand side above as a motivic version of Borel–Moore homology: the cohomology of the dualizing sheaf, in the context of motivic sheaves.

The $A_n(\mathcal{X})$ are just zeroth cohomology groups of the *chain complexes* of derived global sections,

$$C_{\bullet}^{\text{BM}}(\mathcal{X})\langle -n \rangle := R\Gamma(\mathcal{X}, f^! \mathbf{Q}^{\text{mot}}\langle -n \rangle),$$

which we regard as objects of the derived ∞ -category of chain complexes of \mathbf{Q} -vector spaces, where $\langle -n \rangle := (-n)[-2n]$. When \mathcal{X} is a scheme X , $C_{\bullet}^{\text{BM}}(X)\langle -n \rangle$ coincides with the Bloch cycle complex $\mathbf{z}_n(X)$ (see [Kha1, Ex. 2.10]). One may thus think of $C_{\bullet}^{\text{BM}}(\mathcal{X})\langle -n \rangle$ as a Bloch cycle complex for \mathcal{X} .⁷

For the proof of stacky concentration, our key tool is the localization triangle: for any closed substack \mathcal{Z} of \mathcal{X} with open complement $\mathcal{U} = \mathcal{X} \setminus \mathcal{Z}$, there is an exact triangle of complexes

$$C_{\bullet}^{\text{BM}}(\mathcal{Z})\langle -n \rangle \rightarrow C_{\bullet}^{\text{BM}}(\mathcal{X})\langle -n \rangle \rightarrow C_{\bullet}^{\text{BM}}(\mathcal{U})\langle -n \rangle$$

This is a powerful extension of the right-exact localization sequence $A_n(\mathcal{Z}) \rightarrow A_n(\mathcal{X}) \rightarrow A_n(\mathcal{U}) \rightarrow 0$. Using the formalism of higher algebra [Lur], the localization $[\Sigma^{-1}]$ may be performed at the level of complexes, and preserves such exact triangles.

This device allows us to perform a series of reductions. First, to prove that push-forward along a closed immersion induces an isomorphism after Σ -localization, it suffices by localization to demonstrate Σ -acyclicity of motivic Borel–Moore chains on the complement. To show Σ -acyclicity of $C_{\bullet}^{\text{BM}}(\mathcal{X})\langle * \rangle$ for some \mathcal{X} , we may apply localization inductively to a stratification of \mathcal{X} by global quotient stacks to reduce to the case of $\mathcal{X} = [X/\text{GL}_m]$ a global quotient. By a variant of Thomason’s “reduction to the subtorus” argument, which we prove for $C_{\bullet}^{\text{BM}}(-)\langle * \rangle$ (see Theorem 2.5), we can replace GL_m by its maximal subtorus T . This T -action is generically trivial, up to “Morita equivalence”, so yet another application of the localization triangle reduces us to the following straightforward observation: if $\mathcal{X} = [X/G]$ is the quotient of a scheme by the trivial action of a diagonalizable group scheme, such that there exists a line bundle $\mathcal{L} \in \Sigma$ with $c_1(\mathcal{L}|_{BG}) = 0$, then the localization $C_{\bullet}^{\text{BM}}(\mathcal{X})[\Sigma^{-1}]$ vanishes. We refer to Subsect. 3.3 for the details.

0.4. Localization formulas. An immediate consequence of torus concentration is the Atiyah–Bott localization formula, which is an explicit computation of the inverse of the isomorphism i_* . In the situation of Theorem B, suppose X and Z are smooth Artin stacks. Then the inclusion $i : Z \hookrightarrow X$ is lci and we have a Gysin pull-back $i^! : A_*^T(X) \rightarrow A_*^T(Z)$ which satisfies the self-intersection formula

$$i^! \circ i_*(-) = (-) \cap e(N)$$

where $e(N)$ is the top Chern class of the normal bundle. As long as the latter is invertible⁸ in $A_T^*(Z)[\Sigma^{-1}]$, this immediately yields the localization formula

$$(i_*)^{-1} = i^!(-) \cap e(N)^{-1}. \quad (0.6)$$

⁷Note that we do not know of any way to define the Bloch cycle complex $\mathbf{z}_n(\mathcal{X})$ directly (i.e., without passing through motivic cohomology), despite two of the authors’ attempts to do so.

⁸This is true when Z has trivial T -action and finite stabilizers, and N has no T -fixed part; see [AKLPR, Prop. 1.3, Cor. A.31].

Moreover, the same strategy can be adapted to the “virtual” version, where X and Z are *quasi-smooth*. This gives a conceptual proof of the virtual localization formula of [GP] over arbitrary base fields, and shows that it holds without the global resolution and embeddability hypotheses required in *op. cit.* We refer to [AKLPR] for details.

0.5. Contents of the paper. We begin in Sect. 1 by establishing some notation and conventions on algebraic stacks. In Subsect. 1.4 we introduce the G -stabilizer group at a point of an Artin stack X with G -action; note that these are not the same as the stabilizers of the quotient stack $[X/G]$ when X itself has nontrivial stabilizers. Subsect. 1.5 contains a review of various notions of G -fixed loci in the context of stacks, treated in detail in the companion paper [AKLPR].

In Sect. 2 we review the cohomological formalism for intersection theory on stacks developed in [Kha1]. We record the projective bundle formula in this context (Subsect. 2.4). In Subsect. 2.5 we establish an analogue of Thomason’s “reduction to the subtorus”, a very useful technique for reducing statements about G -equivariant intersection theory to the case where G is a split torus.

The stacky concentration theorem (Theorem A) is stated and proven in Sect. 3. We specialize to quotient stacks to derive equivariant concentration (Corollary D) in Subsect. 3.5. Sect. 4 is devoted to the torus-equivariant case (Theorem B), where we can derive a more explicit statement. We end with some examples, including concentration for the moduli stack of Higgs bundles on a curve with its canonical \mathbf{G}_m -scaling action (Corollary C).

0.6. Acknowledgments. We would like to thank Marc Levine and Yan Soibelman for their interest in this work, and Alexandre Minets for help with moduli stacks of Higgs sheaves. We would also like to thank Tasuki Kinjo who participated in this project in the initial stages.

We acknowledge support from the ERC grant QUADAG (D.A.), the NSTC grant 110-2115-M-001-016-MY3 (A.A.K.), the DFG through SFB 1085 Higher Invariants (C.R.), and the EPSRC grant no EP/R014604/1 (A.A.K. and C.R.). We also thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for hospitality during the programme “Algebraic K-theory, motivic cohomology and motivic homotopy theory” where the final revisions of this paper were completed. C.R. is grateful to Max Planck Institute for Mathematics in Bonn for its hospitality and financial support at the time of writing the paper.

1. PRELIMINARIES ON STACKS

1.1. Stacks. We fix a base commutative ring k and denote by Stk_k the ∞ -category of Artin stacks⁹ that are locally of finite type over k and have quasi-compact and separated diagonal. Given an Artin stack $S \in \mathrm{Stk}_k$, we denote by Stk_S the ∞ -category of locally of finite type Artin stacks over S with quasi-compact and separated diagonal.

Let us briefly review the definitions; we refer to [Gai, §4.2] or [Toë, §3.1] for more details. A *prestack* is a presheaf of ∞ -groupoids on the site of k -schemes. A *stack* is a prestack that satisfies hyperdescent with respect to the étale topology.

A stack is *0-Artin* if it is an algebraic space, i.e., if it has schematic and monomorphic (= (-1) -truncated) diagonal¹⁰ and admits a surjective étale morphism from a k -scheme. A stack is *n -Artin*, for $n > 0$, if it has $(n - 1)$ -representable diagonal and admits a surjective smooth morphism from a scheme. A stack is *Artin* if it is n -Artin for some n . A stack is *Deligne–Mumford* if it has representable (= 0-representable) diagonal and admits a surjective étale morphism from a scheme, or equivalently if it is 1-Artin with unramified diagonal.

In the above definitions, a morphism of prestacks $f : X \rightarrow Y$ is called *schematic*, resp. *$(n - 1)$ -representable*, if for every scheme S and every morphism $S \rightarrow Y$, the base change $X \times_Y S$ is a scheme, resp. $(n - 1)$ -Artin. An $(n - 1)$ -representable morphism $f : X \rightarrow Y$ is *étale* (resp. *smooth*, *flat*, *surjective*), if for every scheme S and every morphism $S \rightarrow Y$, the morphism of $(n - 1)$ -representable stacks $X \times_Y S \rightarrow S$ is étale (resp. smooth, flat, surjective).

1.2. Points. A *point* of a prestack X is a field-valued point, i.e., a morphism $x : \mathrm{Spec}(k(x)) \rightarrow X$ where $k(x)$ is a field (which we call the *residue field* at x). A *geometric point* of X is a field-valued point x whose residue field $k(x)$ is an algebraic closure of a residue field of k at a prime ideal \mathfrak{p} . A morphism of stacks is *surjective* if it is surjective on geometric points.

The *set of points* of X , denoted $|X|$, is the colimit

$$\varinjlim_{\kappa} \pi_0 X(\kappa),$$

taken over fields κ , in the category of sets. Here $\pi_0 X(\kappa)$ is the set of connected components of the ∞ -groupoid $X(\kappa)$, and given a field extension $\kappa' \rightarrow \kappa$, the corresponding transition arrow is the map induced by $X(\kappa') \rightarrow X(\kappa)$ on sets of connected components. When X is 1-Artin, $|X|$ admits a canonical structure of topological space (see e.g. [SP, Tag 04XL]).

⁹Note that, except in the introduction, all our stacks are implicitly *higher* by default, i.e., they define sheaves of ∞ -groupoids.

¹⁰hence in particular takes values in sets (= 0-truncated ∞ -groupoids)

1.3. Stabilizers. Let X be a 1-Artin stack and x a point. The *stabilizer* at x is the group algebraic space $\underline{\mathrm{Aut}}_X(x)$ of automorphisms of x . This can be defined equivalently as the fibred product $\mathrm{Spec}(k(x)) \times_X \mathrm{Spec}(k(x))$, the fibre of the diagonal $X \rightarrow X \times X$ over (x, x) , or the fibre of the projection of the inertia stack $I_X \rightarrow X$ over x .

We say a 1-Artin stack X has *affine stabilizers* if for every point x of X , the stabilizer $\underline{\mathrm{Aut}}_X(x)$ is affine. If X has affine inertia or affine diagonal, then it has affine stabilizers.

We say that X has *finite stabilizers* if for every point x the stabilizer $\underline{\mathrm{Aut}}_X(x)$ is finite (over $\mathrm{Spec}(k(x))$), or equivalently if X has quasi-finite diagonal. If X is Deligne–Mumford or has quasi-finite inertia, then it has finite stabilizers.

1.4. Stabilizers of group actions. Given an action of an algebraic group G on an Artin stack X , we define the stabilizer of the action (“ G -stabilizer”) at any point x of X . When the stabilizer at x of X itself is trivial, this coincides with the stabilizer of the quotient stack $[X/G]$ at x .

We fix an fppf group algebraic space G over a k -scheme S .

Remark 1.1. Let $f : X \rightarrow Y$ be a morphism of Artin stacks over S . The relative inertia stack $I_{X/Y}$ is a group Artin stack over X which fits into a cartesian square

$$\begin{array}{ccc} I_{X/Y} & \longrightarrow & I_{X/S} \\ \downarrow & & \downarrow \\ X & \longrightarrow & I_{Y/S} \times_Y X \end{array}$$

of group stacks over X . The lower horizontal arrow is the base change of the identity section $e : Y \rightarrow I_{Y/S}$. When f is representable, $I_{X/Y} \rightarrow X$ is an isomorphism, i.e., $I_{X/S} \rightarrow I_{Y/S} \times_Y X$ is a monomorphism of group stacks. See e.g. [SP, Tag 050P].

Remark 1.2. Let X be an Artin stack over S with G -action and denote by $\mathcal{X} = [X/G]$ the quotient stack. Applying Remark 1.1 to the morphisms $X \rightarrow \mathcal{X}$ and $\mathcal{X} \rightarrow BG$, we get the cartesian squares of group stacks over X

$$\begin{array}{ccccc} X & \longrightarrow & I_{X/S} & \longrightarrow & X \\ \downarrow & & \downarrow & & \downarrow \\ X & \longrightarrow & I_{\mathcal{X}/S} \times_{\mathcal{X}} X & \longrightarrow & G \times_S X, \end{array}$$

where $I_{X/\mathcal{X}} \simeq X$ since $X \rightarrow \mathcal{X}$ is representable and the right-hand vertical arrow is the identity section. For every S -scheme A and every A -valued point x of X , this gives rise to an exact sequence of group algebraic spaces over A

$$1 \rightarrow \underline{\mathrm{Aut}}_X(x) \rightarrow \underline{\mathrm{Aut}}_{\mathcal{X}}(x) \xrightarrow{\alpha_A} G_A \quad (1.3)$$

where $G_A := G \times_S A$ denotes the fibre of G over x .

Definition 1.4. Let X be an Artin stack over S with G -action. For any scheme A and every A -valued point x of X , the G -*stabilizer* (or *stabilizer of*

the G -action) at x is an fppf sheaf of groups $\mathrm{St}_X^G(x)$ defined as the cokernel of the homomorphism $\underline{\mathrm{Aut}}_X(x) \hookrightarrow \underline{\mathrm{Aut}}_{\mathcal{X}}(x)$. Thus we have a short exact sequence

$$1 \rightarrow \underline{\mathrm{Aut}}_X(x) \rightarrow \underline{\mathrm{Aut}}_{\mathcal{X}}(x) \rightarrow \mathrm{St}_X^G(x) \rightarrow 1 \quad (1.5)$$

of sheaves of groups over A . Note that $\mathrm{St}_X^G(x)$ can be regarded as a subgroup of G_A , since it is the image of $\alpha_A : \underline{\mathrm{Aut}}_{\mathcal{X}}(x) \rightarrow G_A$.

Remark 1.6. For a field-valued point $x : \mathrm{Spec}(k(x)) \rightarrow X$, the G -stabilizer $\mathrm{St}_X^G(x)$ is a group algebraic space. This follows from [SGA3, Exp. V, Cor. 10.1.3], since in this case $\underline{\mathrm{Aut}}_X(x)$ is flat over $\mathrm{Spec}(k(x))$. Since X has separated diagonal, $\mathrm{St}_X^G(x)$ is moreover a group *scheme* by [SP, 0B8F].

Remark 1.7. When X has trivial stabilizers (i.e., is an algebraic space), then the G -stabilizer $\mathrm{St}_X^G(x)$ at a point x is nothing else than $\underline{\mathrm{Aut}}_{\mathcal{X}}(x)$, the stabilizer at x of the quotient stack $\mathcal{X} = [X/G]$.

Remark 1.8. Let X be an Artin stack over S with G -action. Let A be an S -scheme and x an A -valued point of X . From the short exact sequence (1.5) we see that the induced morphism $B\underline{\mathrm{Aut}}_X(x) \rightarrow B\underline{\mathrm{Aut}}_{\mathcal{X}}(x)$ is a $\mathrm{St}_X^G(x)$ -torsor, where $\mathcal{X} = [X/G]$. Moreover, there is a commutative diagram

$$\begin{array}{ccc} B\underline{\mathrm{Aut}}_X(x) & \longrightarrow & X \\ \downarrow & & \downarrow \\ B\underline{\mathrm{Aut}}_{\mathcal{X}}(x) & \longrightarrow & \mathcal{X} \end{array}$$

where the right-hand arrow is a G -torsor. In particular, we find that there is a canonical $\mathrm{St}_X^G(x)$ -action on the group algebraic space $\underline{\mathrm{Aut}}_X(x)$, and the canonical monomorphism

$$B\underline{\mathrm{Aut}}_X(x) \hookrightarrow X$$

is equivariant with respect to the $\mathrm{St}_X^G(x)$ -action on the source and G -action on the target.

1.5. Fixed loci. Fix an fppf group algebraic space G over a k -scheme S . Let X be an Artin stack over S with G -action.

Definition 1.9. We define the G -fixed locus $X^G \subseteq X$ as follows. An A -point $x \in X(A)$, where A is an S -scheme, belongs to X^G if and only if the homomorphism of group algebraic spaces over A (1.3)

$$\alpha_A : \underline{\mathrm{Aut}}_{[X/G]}(x) \rightarrow G_A$$

is surjective¹¹. In other words, X^G is the locus of points $x \in X(A)$ where the inclusion of the G -stabilizer $\mathrm{St}_X^G(x) \subseteq G_A$ is an equality (see Definition 1.4).

Definition 1.10. When the subset $|X^G| \subseteq |X|$ is closed, it follows that there exists a reduced closed substack X_{red}^G of X such that $|X_{\mathrm{red}}^G| = |X^G|$. In that case, we refer to $X_{\mathrm{red}}^G \subseteq X$ as the *reduced G -fixed locus*.

¹¹i.e., an effective epimorphism of fppf sheaves: it admits a section after base change along some fppf cover $A' \twoheadrightarrow A$

Definition 1.11. The *homotopy G -fixed point stack* of X , denoted X^{hG} , is defined as follows. An A -point of X^{hG} , where A is an S -scheme, amounts to an A -point $x \in X(A)$ together with a *group-theoretic section* of the homomorphism (1.3)

$$\alpha_A : \underline{\mathrm{Aut}}_{[X/G]}(x) \rightarrow G_A$$

of group algebraic spaces over A . The projection $X^{hG} \rightarrow X$ factors through the fixed locus $X^G \subseteq X$ by definition.

We recall the following results from [AKLPR, App. A]:

Theorem 1.12. *Let S be a k -scheme and $G \rightarrow S$ an fppf group algebraic space with connected fibres. Then for every Artin stack X over S with G -action, we have:*

- (i) *Suppose X is an algebraic space. Then there is a canonical isomorphism $X^G \simeq X_{\mathrm{cl}}^{hG}$ over X . Moreover, the morphisms $X^G \rightarrow X$ and $X^{hG} \rightarrow X$ are closed immersions.*
- (ii) *Suppose X is locally of finite type over k and 1-Artin with finite stabilizers. Then the subset $|X^G| \subseteq |X|$ is closed. In particular, the reduced G -fixed locus $X_{\mathrm{red}}^G \subseteq X$ exists.*
- (iii) *Suppose S is locally noetherian, $G = T \rightarrow S$ is a split torus, and X is of finite type over k and 1-Artin with affine stabilizers. Then there exists an isogeny $\rho : T' \rightarrow T$ where T' is a split torus over S and a canonical surjection $X_{\mathrm{red}}^{hT'} \rightarrow X_{\mathrm{red}}^T$ over X , where the target is the reduced T -fixed locus. In particular, the set-theoretic image of $X^{hT'} \rightarrow X$ coincides with $|X^T| \subseteq |X|$.*
- (iv) *Suppose S is locally noetherian, $G = T \rightarrow S$ is a split torus, and X is locally of finite type over k and Deligne–Mumford with quasi-compact separated diagonal. Then the canonical morphism $X^{hT} \rightarrow X$ is a closed immersion.*

Proof. See Prop. A.21, Prop. A.12, Cor. A.42, and Prop. A.23, respectively, of [AKLPR]. \square

2. INTERSECTION THEORY ON STACKS

In this section we briefly summarize the definitions and main properties of motivic Borel–Moore homology of Artin stacks from [Kha1].

2.1. Motivic Borel–Moore homology.

2.1.1. *Motivic spectra.* Let $\mathbf{SH}(k)$ denote the stable ∞ -category of motivic spectra over $\mathrm{Spec}(k)$; see e.g. [Voe, Kha2] for definitions. We let $\Lambda \in \mathbf{SH}(k)$

be a motivic commutative ring spectrum which is \mathbf{Q} -linear and oriented, such as

$$\begin{aligned} \Lambda &= \mathbf{Q}^{\text{mot}} && \text{(motivic cohomology),} \\ \Lambda &= \text{KGL}_{\mathbf{Q}} && \text{(algebraic K-theory),} \\ \Lambda &= \text{MGL}_{\mathbf{Q}} && \text{(algebraic cobordism).} \end{aligned}$$

see Examples 2.10, 2.12, and 2.13 of [Kha1]. By [CD, Cor. 14.2.16], we may (and will) regard Λ as a \mathbf{Q}^{mot} -algebra.

We consider the twists

$$\Lambda\langle n \rangle := \Lambda(n)[2n] \in \mathbf{SH}(k)$$

where $\Lambda(n)$ is the Tate twist by $n \in \mathbf{Z}$ (see [CD, Def. 2.4.17]).

2.1.2. Generalized Borel–Moore chains. Given an Artin stack $X \in \text{Stk}_k$, we have the complexes of *Borel–Moore chains* with coefficients in Λ (see [Kha1, §2.1]):

$$\mathbf{C}_{\bullet}^{\text{BM}}(X; \Lambda) := R\Gamma(X, a_X^! \Lambda),$$

where $a_X : X \rightarrow \text{Spec}(k)$ is the structural morphism and $a_X^!$ is the $!$ -inverse image functor constructed in [Kha1, Thm. A.5]. We regard this as a *complex*, which here is an abbreviation of “object of the derived ∞ -category $\mathbf{D}(\mathbf{Q})$ of chain complexes of \mathbf{Q} -vector spaces”.

We will often omit the coefficient when it is understood, and write

$$\mathbf{C}_{\bullet}^{\text{BM}}(X)\langle n \rangle := \mathbf{C}_{\bullet}^{\text{BM}}(X; \Lambda\langle n \rangle)$$

for the twists.

Explicitly, $\mathbf{C}_{\bullet}^{\text{BM}}(X)$ can be described as a homotopy limit

$$\mathbf{C}_{\bullet}^{\text{BM}}(X) \simeq \varprojlim_{(U, u)} \mathbf{C}_{\bullet}^{\text{BM}}(U)\langle -d_u \rangle$$

over pairs (U, u) where $U \in \text{Stk}_k$ is a scheme and $u : U \rightarrow X$ is a smooth morphism of relative dimension d_u .

When X is a global quotient stack $[U/G]$, $\mathbf{C}_{\bullet}^{\text{BM}}(X)$ can be described as Borel-type G -equivariant Borel–Moore chains on U ; see [KR1, Cor. 6.2].

2.1.3. Operations. Each $\mathbf{C}_{\bullet}^{\text{BM}}(X; \Lambda)\langle n \rangle$ is a module over the complex of cochains,

$$\mathbf{C}^{\bullet}(X; \Lambda) := R\Gamma(X, a_X^* \Lambda)$$

by cap product (see [Kha1, 2.2.6]).

Given a proper morphism $f : X \rightarrow Y$ in Stk_k , there is a push-forward map

$$f_* : \mathbf{C}_{\bullet}^{\text{BM}}(X) \rightarrow \mathbf{C}_{\bullet}^{\text{BM}}(Y)$$

induced by the counit $f_* f^! \simeq f_! f^! \rightarrow \text{id}$ (see [Kha1, 2.2.1]).

Given a smooth morphism $f : X \rightarrow Y$ of relative dimension d in Stk_k , there is a Gysin pull-back map

$$f^! : C_{\bullet}^{\text{BM}}(Y) \rightarrow C_{\bullet}^{\text{BM}}(X)\langle -d \rangle$$

induced by the cotrace $\text{id} \rightarrow f_* f^! \langle -d \rangle$ (see [Kha1, 2.2.2]).

2.1.4. *Relative and equivariant versions.* For a morphism $f : X \rightarrow S$ in Stk_k , we consider *relative Borel–Moore chains*:

$$C_{\bullet}^{\text{BM}}(X/S; \Lambda) := R\Gamma(X, f^!(\Lambda_S)),$$

where $\Lambda_S = a_S^*(\Lambda)$ for $a_S : S \rightarrow \text{Spec}(k)$ the projection. As in the absolute case, these admit proper push-forward and smooth Gysin pull-back functoriality in X . Note that $C_{\bullet}^{\text{BM}}(X) = C_{\bullet}^{\text{BM}}(X/k)$ and $C^{\bullet}(X) = C_{\bullet}^{\text{BM}}(X/X)$.

Let G be an fppf group algebraic space over k . For every $X \in \text{Stk}_k$ with G -action, we write

$$\begin{aligned} C_G^{\bullet}(X) &:= C^{\bullet}([X/G]), \\ C_{\bullet}^{\text{BM},G}(X) &:= C_{\bullet}^{\text{BM}}([X/G]_{/BG}), \end{aligned}$$

for G -equivariant cochains and G -equivariant Borel–Moore chains on X . Since BG is smooth of dimension $-\dim(G)$, we have

$$C_{\bullet}^{\text{BM},G}(X) \simeq C_{\bullet}^{\text{BM}}([X/G])\langle -\dim(G) \rangle.$$

2.1.5. *Periodization.* We consider the commutative Λ -algebra

$$\Lambda\langle * \rangle := \Lambda[\gamma, \gamma^{-1}] \simeq \Lambda \otimes_{\mathbf{Q}^{\text{mot}}} \mathbf{Q}^{\text{mot}}[\gamma, \gamma^{-1}] \in \mathbf{SH}(k)$$

where $\gamma : \Lambda\langle 1 \rangle \rightarrow \Lambda\langle * \rangle$ is a free invertible generator (cf. [CD, Cor. 14.2.17(3)]). At the level of Λ -modules, we have $\Lambda\langle * \rangle \simeq \bigoplus_{i \in \mathbf{Z}} \Lambda\langle i \rangle$. Note that $\Lambda \mapsto \Lambda\langle * \rangle$ commutes with $*$ -pullback along $X \rightarrow \text{Spec}(k)$.

We also consider the $C^{\bullet}(X)$ -algebras

$$\begin{aligned} C^{\bullet}(X)\langle * \rangle &:= C^{\bullet}(X)[\gamma, \gamma^{-1}] := \bigoplus_{i \in \mathbf{Z}} C^{\bullet}(X)\langle i \rangle, \\ \widehat{C}^{\bullet}(X)\langle * \rangle &:= C^{\bullet}(X)[[\gamma, \gamma^{-1}]] := \prod_{i \in \mathbf{Z}} C^{\bullet}(X)\langle i \rangle. \end{aligned}$$

There are canonical maps

$$C^{\bullet}(X)\langle * \rangle \rightarrow C^{\bullet}(X; \Lambda\langle * \rangle) \rightarrow \widehat{C}^{\bullet}(X)\langle * \rangle. \quad (2.1)$$

We consider the periodized Borel–Moore chain complexes

$$\begin{aligned} C_{\bullet}^{\text{BM}}(X)\langle * \rangle &:= C_{\bullet}^{\text{BM}}(X)[\gamma, \gamma^{-1}], \\ \widehat{C}_{\bullet}^{\text{BM}}(X)\langle * \rangle &:= C_{\bullet}^{\text{BM}}(X)[[\gamma, \gamma^{-1}]] \end{aligned}$$

as modules over $C^{\bullet}(X)\langle * \rangle$ and $\widehat{C}^{\bullet}(X)\langle * \rangle$, respectively.

For G an fppf group algebraic space over k , we define $C_G^{\bullet}(X)\langle * \rangle$, $\widehat{C}_G^{\bullet}(X)\langle * \rangle$, $C_{\bullet}^{\text{BM},G}(X)\langle * \rangle$, and $\widehat{C}_{\bullet}^{\text{BM},G}(X)\langle * \rangle$ similarly.

2.1.6. *Chern character.* The Chern character induces a canonical isomorphism

$$\mathrm{KGL}_{\mathbf{Q}} \simeq \mathbf{Q}^{\mathrm{mot}}\langle * \rangle$$

of $\mathbf{Q}^{\mathrm{mot}}$ -algebras in $\mathbf{SH}(k)$ by [CD, Cor. 14.2.17(3)]. Via the $\mathbf{Q}^{\mathrm{mot}}\langle * \rangle$ -algebra structure on $\Lambda\langle * \rangle$ we obtain a canonical map

$$\mathrm{ch} : \mathrm{KGL}_{\mathbf{Q}} \rightarrow \Lambda\langle * \rangle$$

of $\mathbf{Q}^{\mathrm{mot}}$ -algebras.

For every $X \in \mathrm{Stk}_k$ this yields a canonical map

$$\mathrm{C}^{\bullet}(X; \mathrm{KGL}_{\mathbf{Q}}) \rightarrow \mathrm{C}^{\bullet}(X; \Lambda\langle * \rangle) \rightarrow \widehat{\mathrm{C}}^{\bullet}(X; \Lambda)\langle * \rangle$$

where the second arrow is as in (2.1). Recall from [Kha1, Ex. 2.13] that $\mathrm{C}^{\bullet}(X; \mathrm{KGL}_{\mathbf{Q}})$ is identified with the étale K-theory of X (see [Kha3, §5.2]). Via the canonical map from K-theory to its étale localization, we thus get a Chern character map

$$\mathrm{ch} : \mathrm{K}(X) \rightarrow \mathrm{K}^{\mathrm{ét}}(X)_{\mathbf{Q}} \simeq \mathrm{C}^{\bullet}(X; \mathrm{KGL}_{\mathbf{Q}}) \rightarrow \widehat{\mathrm{C}}^{\bullet}(X; \Lambda)\langle * \rangle.$$

In particular, K-theory classes $\alpha \in \mathrm{K}(X)$ act on $\widehat{\mathrm{C}}_{\bullet}^{\mathrm{BM}}(X; \Lambda)\langle * \rangle$.

For example, for any fppf group algebraic space G over k , there is a ring homomorphism

$$\mathrm{R}(G) \simeq \mathrm{K}_0(BG) \rightarrow \pi_0 \widehat{\mathrm{C}}^{\bullet}(BG; \Lambda)\langle * \rangle$$

where $\mathrm{R}(G)$ is the representation ring of G . When G acts on $X \in \mathrm{Stk}_k$, $\pi_0 \widehat{\mathrm{C}}_{\bullet}^{\mathrm{BM}, G}(X; \Lambda)$ is thus a $\mathrm{R}(BG)$ -module.

2.2. Chow, G-theory, bordism.

2.2.1. *Chow homology.* For $\Lambda = \mathbf{Q}^{\mathrm{mot}}$ the motivic cohomology spectrum, we write

$$\mathrm{A}_n(X, s) := \pi_s \mathrm{C}_{\bullet}^{\mathrm{BM}}(X; \mathbf{Q}^{\mathrm{mot}}\langle -n \rangle) \simeq \mathrm{H}^{-2n-s}(X; a_X^! \mathbf{Q}^{\mathrm{mot}}\langle -n \rangle)$$

for every $X \in \mathrm{Stk}_k$ and $n, s \in \mathbf{Z}$, and $\mathrm{A}_n(X) := \mathrm{A}_n(X, 0)$.

Suppose X is a global quotient stack, e.g. $[U/G]$ where U is an algebraic space of finite type over k and G is a linear algebraic group over k acting on U . Then $\mathrm{A}_n(X)$ is identified with the equivariant Chow group $\mathrm{A}_{n-\dim(G)}^G(U) \otimes \mathbf{Q}$ of [EG1] by [KR1, Cor. 6.5]. In particular, it is identified with Kresch's Chow group [Kre] in this case.

Suppose X is a 1-Artin stack with finite inertia¹². Then X admits a coarse moduli space $\pi : X \rightarrow M$ [KM] and the push-forward $\pi_* : \mathrm{A}_n(X) \rightarrow \mathrm{A}_n(M)$ is an isomorphism by proper codescent (see [Kha3, Thm. 5.7]). In particular, $\mathrm{A}_*(X)$ is identified with Kresch's Chow groups also in this case.

In fact, for any 1-Artin stack X there is a canonical map to $\mathrm{A}_n(X)$ from Kresch's Chow group, specializing to the above isomorphisms. It will be

¹²When X is separated, this is equivalent to having finite stabilizers.

shown in [BP] that it is invertible as long as X has affine stabilizers (and hence admits a stratification by global quotient stacks).

2.2.2. *G-theory.* For $\Lambda = \mathrm{KGL}_{\mathbf{Q}}$ the algebraic K-theory spectrum, we write

$$G_s^{\acute{e}t}(X) := \pi_s C_{\bullet}^{\mathrm{BM}}(X; \mathrm{KGL}_{\mathbf{Q}})$$

for every $X \in \mathrm{Stk}_k$ and $s \in \mathbf{Z}$. These are the étale G-theory groups of X (see [Kha1, Ex. 2.13] and [Kha3, §5.2]). Recall that we have $\mathrm{KGL}_{\mathbf{Q}}\langle n \rangle \simeq \mathrm{KGL}_{\mathbf{Q}}$ for all $n \in \mathbf{Z}$. We have the étale-local Grothendieck–Riemann–Roch isomorphism

$$G_s^{\acute{e}t}(X) \simeq \prod_{n \in \mathbf{Z}} A_n(X, s)$$

for all s by [Kha1, §3.5].

For $X = [U/G]$ a quotient stack, $G_s^{\acute{e}t}(X)$ is Borel-equivariant G-theory $G_s^{\acute{e}t, G}(U)_{\mathbf{Q}}$ by [KR1, Cor. 7.8]. In particular, $G_0^{\acute{e}t}(X)$ is the *completion* $G_0^G(U)_{I_G}^{\wedge}$ of the Grothendieck group of G -equivariant coherent sheaves at the augmentation ideal I_G .

2.2.3. *Bordism.* For $\Lambda = \mathrm{MGL}_{\mathbf{Q}}$ the algebraic cobordism spectrum, we write

$$\Omega_n(X, s) := \pi_s C_{\bullet}^{\mathrm{BM}}(X; \mathrm{MGL}_{\mathbf{Q}}\langle -n \rangle)$$

for every $X \in \mathrm{Stk}_k$ and $n, s \in \mathbf{Z}$, and $\Omega_n(X) := \Omega_n(X, 0)$.

When $X = [U/G]$ is a quotient stack and k is of characteristic zero, $\Omega_n(X)$ is identified with the equivariant bordism group $\Omega_{n-\dim(G)}^G(U) \otimes \mathbf{Q}$ of [HML] by [KR1, Thm. 8.4].

2.2.4. *Betti/ ℓ -adic Borel–Moore homology.* The definition of Borel–Moore chains makes sense in any appropriate six functor formalism. Specifically, for the purposes of this paper, one can replace $\mathbf{SH}(-)$ by any étale-local oriented topological weave [Kha4]; see also [KR2].

For example, over $k = \mathbf{C}$ we may take $\Lambda = \mathbf{Q}$ in the derived ∞ -category of abelian groups, and form $C_{\bullet}^{\mathrm{BM}}(X; \mathbf{Q})$ using the $!$ -pullback functor on derived ∞ -categories of sheaves of abelian groups. See e.g. [KK, §1.1] for the six functor formalism on Artin stacks in this context.

Similarly, over a field k in which ℓ is invertible, we may form $C_{\bullet}^{\mathrm{BM}}(X; \mathbf{Q}_{\ell})$ using the six functor formalism for ℓ -adic sheaves on Artin stacks [LZ1, LZ2].

2.3. Localization triangle.

Theorem 2.2. *Let $S \in \mathrm{Stk}_k$, $X \in \mathrm{Stk}_S$, and $i : Z \rightarrow X$, $j : U \rightarrow X$ a pair of complementary closed and open immersions. Then there is a canonical exact triangle*

$$C_{\bullet}^{\mathrm{BM}}(Z/S) \xrightarrow{i_*} C_{\bullet}^{\mathrm{BM}}(X/S) \xrightarrow{j^!} C_{\bullet}^{\mathrm{BM}}(U/S)$$

in $\mathbf{D}(S)$.

Proof. By [Kha1, Thm. 2.18] we have for every motivic sheaf \mathcal{F} on X the exact triangle

$$i_* i^! (\mathcal{F}) \rightarrow \mathcal{F} \rightarrow j_* j^! (\mathcal{F}).$$

The claim follows by taking $\mathcal{F} = f^! (\Lambda_S)$, where $f : X \rightarrow S$ is the structural morphism, and applying derived global sections. \square

2.4. Projective bundle formula.

Theorem 2.3. *Let $S \in \text{Stk}_k$ and $X \in \text{Stk}_S$. Let \mathcal{E} be a locally free sheaf on X of rank $r + 1$ ($r \geq 0$), and write $\pi : P = \mathbf{P}(\mathcal{E}) \rightarrow X$ for the associated projective bundle. Then the maps*

$$\pi^! (-) \cap e(\mathcal{O}(-1))^{\cup i} : C_{\bullet}^{\text{BM}}(X/S) \rightarrow C_{\bullet}^{\text{BM}}(P/S)\langle -r + i \rangle,$$

induce a canonical isomorphism of $C^{\bullet}(X)$ -modules

$$\bigoplus_{0 \leq i \leq r} C_{\bullet}^{\text{BM}}(X/S)\langle r - i \rangle \rightarrow C_{\bullet}^{\text{BM}}(P/S). \quad (2.4)$$

Proof. Since $C_{\bullet}^{\text{BM}}(-/S)$ satisfies étale descent, the map (2.4) is the limit over smooth morphisms $t : T \rightarrow X$ with T a scheme of the analogous maps

$$\bigoplus_{0 \leq i \leq r} C_{\bullet}^{\text{BM}}(T/S)\langle r - i - d_t \rangle \rightarrow C_{\bullet}^{\text{BM}}(T \times_S P/S)\langle -d_t \rangle,$$

where d_t is the relative dimension of $t : T \rightarrow X$. We may therefore assume that X is a scheme. By Zariski descent on X , we may further assume that \mathcal{E} admits a trivial direct summand. It is therefore enough to show the following:

- (*) If (2.4) is invertible for $\mathcal{E} = \mathcal{E}_0$ (locally free of rank r), then it is also invertible for $\mathcal{E} = \mathcal{E}_0 \oplus \mathcal{O}$.

Let $P_0 = \mathbf{P}(\mathcal{E}_0)$, $P = \mathbf{P}(\mathcal{E}_0 \oplus \mathcal{O})$, and $E = \mathbf{V}(\mathcal{E}) = P \setminus P_0$. Consider the following diagram:

$$\begin{array}{ccccc} \bigoplus_0^{r-1} C_{\bullet}^{\text{BM}}(X/S)\langle r - 1 - i \rangle & \longrightarrow & \bigoplus_0^r C_{\bullet}^{\text{BM}}(X/S)\langle r - i \rangle & \longrightarrow & C_{\bullet}^{\text{BM}}(X/S)\langle r \rangle \\ \downarrow & & \downarrow & & \downarrow \\ C_{\bullet}^{\text{BM}}(P_0/S) & \longrightarrow & C_{\bullet}^{\text{BM}}(P/S) & \longrightarrow & C_{\bullet}^{\text{BM}}(E/S) \end{array}$$

where the upper row is split exact and the lower horizontal row is the localization triangle. The right-hand vertical arrow is Gysin along the projection $E \rightarrow X$ (which is invertible by homotopy invariance), and the left-hand and middle vertical arrows are (2.4) for $\mathcal{E} = \mathcal{E}_0$ and $\mathcal{E} = \mathcal{E}_0 \oplus \mathcal{O}$, respectively. Hence it will suffice to show that this diagram is commutative.

The inclusion $i : P_0 \hookrightarrow P$ exhibits P_0 as the zero locus on P of the cosection $\pi^*(\mathcal{O}_P(-1)) \oplus \mathcal{O}_P \rightarrow \mathcal{O}_P$ (the projection onto the second component), so we have a canonical homotopy

$$i_* i^! (-) \simeq (-) \cap e(\mathcal{O}_P(-1))$$

of maps $C_{\bullet}^{\text{BM}}(P/S) \rightarrow C_{\bullet}^{\text{BM}}(P/S)\langle 1 \rangle$. The induced homotopy

$$i_* \pi_0^! (-) \simeq \pi^! (-) \cap e(\mathcal{O}_P(-1)),$$

where $\pi_0 : P_0 \rightarrow X$ and $\pi : P \rightarrow X$ are the projections, then gives rise to a homotopy up to which the left-hand square in the above diagram commutes. Since $\mathcal{O}_P(-1)$ on P restricts to the trivial line bundle on E , there are canonical homotopies $e(\mathcal{O}_P(-1))^{\cup i}|_E \simeq 0$ for all $i > 0$, whence a homotopy up to which the right-hand square also commutes. The claim follows. \square

2.5. Reduction to the subtorus. Note that the first part of the statement below holds for Λ not necessarily \mathbf{Q} -linear.

Theorem 2.5. *Let $S \in \text{Stk}_k$ and $X \in \text{Stk}_S$. Suppose $G = \text{GL}_n$ acts on X and denote by $T \subseteq G$ the subgroup of diagonal matrices. Then Gysin pull-back along the smooth morphism $p : [X/T] \rightarrow [X/G]$*

$$p^! : C_{\bullet}^{\text{BM}}([X/G]_{/S}) \rightarrow C_{\bullet}^{\text{BM}}([X/T]_{/S})\langle -d \rangle,$$

where $d = \dim(G/T)$, admits a $C^{\bullet}([X/G])$ -module retract. Moreover, it induces an isomorphism

$$p^! : C_{\bullet}^{\text{BM}}([X/G]_{/S})b\mathbf{Q} \rightarrow C_{\bullet}^{\text{BM}}([X/T]_{/S})^{hW},$$

where $(-)^{hW}$ denotes homotopy invariants with respect to the W -action induced by the canonical W -action on $[X/T]$.

Proof. Since $[X/T] \rightarrow [X/B]$ is an affine bundle, where $B \subseteq G$ is the Borel subgroup, the Gysin map

$$C_{\bullet}^{\text{BM}}([X/B]_{/S}) \rightarrow C_{\bullet}^{\text{BM}}([X/T]_{/S})$$

is invertible by homotopy invariance. Therefore, it is enough to show that the Gysin map

$$C_{\bullet}^{\text{BM}}([X/G]_{/S}) \rightarrow C_{\bullet}^{\text{BM}}([X/B]_{/S})$$

admits a retract. Under the Morita isomorphism $[X/B] \simeq [(X \times G/B)/G]$ (with G acting on G/B by conjugation), this is identified with the Gysin map

$$f^! : C_{\bullet}^{\text{BM}}([X/G]_{/S}) \rightarrow C_{\bullet}^{\text{BM}}([(X \times G/B)/G]_{/S})$$

where $f : [(X \times G/B)/G] \rightarrow [X/G]$ is the canonical morphism. Note that f is proper (since G/B is proper over k), so that there is a proper push-forward f_* . We claim that this provides the desired retract. Indeed, since G/B is the scheme of complete flags in \mathbf{A}_X^n over X , the projection f factors as a sequence of iterated projective bundles. Thus the claim follows from the fact that for a projective bundle π , the pullback $\pi^!$ admits a retract by the projective bundle formula (Theorem 2.3), cf. [Ful, Ex. 3.3.5]. The second part follows similarly from the projective bundle formula; see again *loc. cit.* and also recall that with rational coefficients, the homotopy groups of the homotopy invariants can be computed as the ordinary invariants of the homotopy groups. \square

3. CONCENTRATION

In this section we assume the base ring k is noetherian. Let $\Lambda \in \mathbf{D}(k)$ be an object, fixed throughout the section. As per our conventions recalled in Subsect. 2.1, we will leave Λ implicit in the notation.

3.1. The master theorem. Given any set Σ of line bundles on an Artin stack \mathcal{X} , our main result is a stabilizer-wise criterion for torsionness of Borel–Moore chains on \mathcal{X} with respect to the first Chern classes of the line bundles in Σ .

Notation 3.1. Let $\mathcal{X} \in \text{Stk}_k$ be quasi-compact and let $\Sigma \subseteq \text{Pic}(\mathcal{X})$ be a subset. We consider the localizations

$$C^\bullet(\mathcal{X})_{\Sigma\text{-loc}} := C^\bullet(\mathcal{X})(*)[c_1(\Sigma)^{-1}]$$

and

$$C_\bullet^{\text{BM}}(\mathcal{X})_{\Sigma\text{-loc}} := C_\bullet^{\text{BM}}(\mathcal{X})(*)[c_1(\Sigma)^{-1}] \simeq C_\bullet^{\text{BM}}(\mathcal{X}) \otimes_{C^\bullet(\mathcal{X})} C^\bullet(\mathcal{X})_{\Sigma\text{-loc}}.$$

We omit Σ from the subscript when there is no risk of confusion.

Remark 3.2. On homotopy groups, we have

$$\pi_s C_\bullet^{\text{BM}}(\mathcal{X})_{\Sigma\text{-loc}} \simeq \pi_s(C_\bullet^{\text{BM}}(\mathcal{X})(*)[c_1(\Sigma)^{-1}])$$

by [Lur, Prop. 7.2.3.25(2)]. In particular, for $\Lambda = \mathbf{Q}^{\text{mot}}$ and $s = 0$ we recover the localization $A_*(\mathcal{X})_{\text{loc}}$ appearing in the statement of Theorem A.

Definition 3.3. Let $\mathcal{X} \in \text{Stk}_k$ and $\Sigma \subseteq \text{Pic}(\mathcal{X})$ a subset. We introduce the following condition on Σ :

(L) For every geometric point x of \mathcal{X} , there exists an invertible sheaf $\mathcal{L}(x) \in \Sigma$ whose restriction $\mathcal{L}(x)|_{B\text{Aut}_{\mathcal{X}}(x)}$ is trivial.

Remark 3.4. If k is a field, we may also consider variants of condition (L) where we look at k -rational points or \bar{k} -rational points for an algebraic closure \bar{k} of k . The distinction between these versions will play no role in our proofs, so we may sometimes abuse language by referring to any of them as “condition (L)” when there is no risk of confusion. The same applies for other variants of this condition that we will consider below, such as conditions (K) and (L_G).

The main result of this section is as follows:

Theorem 3.5. *Let $S \in \text{Stk}_k$, $\mathcal{X} \in \text{Stk}_S$ quasi-compact 1-Artin with affine stabilizers, and $\Sigma \subseteq \text{Pic}(\mathcal{X})$ a subset. If Σ satisfies condition (L) for \mathcal{X} , then we have*

$$C_\bullet^{\text{BM}}(\mathcal{X}/S)_{\Sigma\text{-loc}} = 0.$$

Corollary 3.6. *Let $\mathcal{X} \in \text{Stk}_k$ be quasi-compact 1-Artin with affine stabilizers. If Σ satisfies condition (L) for \mathcal{X} , then we have $C^\bullet(\mathcal{X})_{\Sigma\text{-loc}} = 0$.*

Proof. Take $S = \mathcal{X}$ in Theorem 3.5. □

Corollary 3.7. *Let $S \in \text{Stk}_k$, $\mathcal{Z}, \mathcal{X} \in \text{Stk}_k$ quasi-compact 1-Artin with affine stabilizers, and $i : \mathcal{Z} \rightarrow \mathcal{X}$ a closed immersion over S . Let $\Sigma \subseteq \text{Pic}(\mathcal{X})$ be a subset satisfying condition (L) for $\mathcal{X} \setminus \mathcal{Z}$. Then the direct image map*

$$i_* : C_\bullet^{\text{BM}}(\mathcal{Z}/S)_{\Sigma\text{-loc}} \rightarrow C_\bullet^{\text{BM}}(\mathcal{X}/S)_{\Sigma\text{-loc}}$$

is invertible.

Proof. Follows by the localization triangle. \square

We will also consider the following variant where Σ is a set of K-theory classes.

Variante 3.8. Let $\Sigma \subseteq K_0(\mathcal{X})$ be a set of *K-theory* classes on \mathcal{X} . We may consider the following variant of condition **(L)**:

(K) For every geometric point x of \mathcal{X} , we have

$$K_0(B\text{Aut}_{\mathcal{X}}(x))[\Sigma^{-1}] = 0,$$

where the action of Σ is given by inverse image along $B\text{Aut}_{\mathcal{X}}(x) \rightarrow \mathcal{X}$.

Remark 3.9. Condition **(K)** can be reformulated as follows: for every geometric point x of \mathcal{X} , there exists a K-theory class $\alpha(x) \in K_0(\mathcal{X})$ which belongs to the multiplicative closure of Σ such that $\alpha(x)|_{B\text{Aut}_{\mathcal{X}}(x)} = 0$.

Recall from (2.1.6) that there is a canonical action of K-theory on Borel–Moore chains via the Chern character. We consider the localizations at $\text{ch}(\Sigma) \subseteq \pi_0 \widehat{C}^\bullet(\mathcal{X})\langle * \rangle$:

$$\begin{aligned} \widehat{C}^\bullet(\mathcal{X})_{\Sigma\text{-loc}} &:= \widehat{C}^\bullet(\mathcal{X})\langle * \rangle[\text{ch}(\Sigma)^{-1}] \\ \widehat{C}_\bullet^{\text{BM}}(\mathcal{X})_{\Sigma\text{-loc}} &:= \widehat{C}_\bullet^{\text{BM}}(\mathcal{X})\langle * \rangle[\text{ch}(\Sigma)^{-1}]. \end{aligned}$$

We have the following analogue of Theorem 3.5:

Theorem 3.10. *Let $S \in \text{Stk}_k$, $\mathcal{X} \in \text{Stk}_S$ quasi-compact 1-Artin with affine stabilizers, and $\Sigma \subseteq K_0(\mathcal{X})$ a subset. If Σ satisfies condition **(K)** for \mathcal{X} , then we have*

$$\widehat{C}_\bullet^{\text{BM}}(\mathcal{X}/S)_{\Sigma\text{-loc}} = 0.$$

3.2. Generic slices. The proof of Theorem 3.5 will require a useful result of Thomason (see [Tho1, Thm. 4.10, Rem. 4.11]) which we reformulate here in stacky terms for the reader's convenience.

Proposition 3.11. *Let T be a diagonalizable torus of finite type over a noetherian scheme S . Let X be a reduced algebraic space of finite type over S with T -action. Then there exists a T -invariant nonempty affine open subspace $U \subseteq X$ such that there is a canonical isomorphism*

$$[U/T] \simeq BT' \times_S V$$

of stacks over BT , for some subgroup $T' \subseteq T$ and affine scheme V over S .

Proof. Recall that the schematic locus $X_0 \subseteq X$ is a nonempty open (see [SP, Tags 03JH, 03JG]). We claim that X_0 is T -invariant, i.e., the action morphism $T \times_S X_0 \rightarrow X$ factors through X_0 . In other words, the open immersion $T \times_S X_0 \times_X X_0 \rightarrow T \times_S X_0$ (base change of the inclusion $X_0 \rightarrow X$) is invertible. We may assume that S is the spectrum of an algebraically closed field k , and it will suffice to check this on k -points. Let $t : \text{Spec}(k) \rightarrow T$ be a k -point. Then t defines an isomorphism $t : X \rightarrow X$. Then $t(X_0) \subseteq X$ is a scheme and hence $t(X_0) \subseteq X_0$ (since X_0 is the largest open which is a scheme). Since t was arbitrary, this shows that X_0 is T -invariant.

Replacing X by X_0 , we may assume that X is a scheme. Now the claim follows by combining [Tho1, Lem. 4.3]¹³ and [Tho1, Thm. 4.10, Rem. 4.11]. \square

3.3. Proof of Theorem 3.5. Let the notation be as in Theorem 3.5. By nil-invariance of Borel–Moore homology, we may replace all stacks by their reductions. Since \mathcal{X} has affine stabilizers, it admits a stratification by global quotient stacks (see [HR, Prop. 2.6]), so again using the localization triangle we may assume that $\mathcal{X} = [X/G]$, where X is a reduced quasi-affine scheme of finite type over k and $G = \mathrm{GL}_n$, $n \geq 0$. Using Theorem 2.5, we may replace G by its maximal subtorus $\mathbf{G}_m^{\times n}$ and thereby assume G is a split torus. (Note that the condition on Σ is clearly preserved under this change of notation.)

By Proposition 3.11, there is a nonempty G -invariant affine open U of X and a diagonalizable subgroup H of G such that $[U/G] \simeq Y \times BH$ with Y an affine scheme. By noetherian induction and the localization triangle, we may replace X by U , and then further replace X by Y and G by H so that $\mathcal{X} = X \times BG$. (Again, note that the condition on Σ is preserved.)

We can assume X is nonempty, so that by condition (L) there exists a geometric point x of X and an invertible sheaf $\mathcal{L} \in \mathrm{Pic}(X)$ such that $\mathcal{L}|_{BG_{k(x)}}$ is trivial. It is enough to show that $c = c_1(\mathcal{L})$ is nilpotent as an element of the ring $\pi_0 \mathbf{C}^\bullet(\mathcal{X})(*)$.

Since G is diagonalizable, we may write $\mathcal{L} = \mathcal{L}_m|_X \otimes m\mathcal{O}_{BG}|_X$ where m is a character of G and \mathcal{L}_m is an invertible sheaf on X (see [SGA3, Exp. I, 4.7.3]). The equality $c|_{BG_{k(x)}} = 0$ thus implies that the invertible sheaf $\mathcal{L}|_{BG_{k(x)}}$ is trivial, i.e., $m = 1$ and $\mathcal{L} \simeq \mathcal{L}_1|_X$. Thus $c = c_1(\mathcal{L}_1)|_X$. Since X is a scheme, $c_1(\mathcal{L}_m)$ is nilpotent (see e.g. [Dég, Prop. 2.1.22(1)]), hence so is c .

Remark 3.12. One can always find a *finite* subset $\Sigma_0 \subseteq \Sigma$ such that the conclusion of Theorem 3.5 still holds. Indeed, note that after the reductions we only need a *single* element of Σ , and in each reduction step we only need to use finitely many more elements.

3.4. Proof of Theorem 3.10. After the same reductions as in the proof of Theorem 3.5, we reduce to the case where $\mathcal{X} = X \times BG$ with X a (nonempty) reduced quasi-affine scheme and G a diagonalizable group scheme. By condition (K), there exists a geometric point x of \mathcal{X} and a class $\alpha \in \Sigma$ with $\alpha|_{BG_{k(x)}}$ trivial, where $G_{k(x)} \simeq \underline{\mathrm{Aut}}_{\mathcal{X}}(x)$ is the base change of G to the residue field $k(x)$. It is enough to show that α is nilpotent as an element of $\mathbf{K}_0(\mathcal{X})$. If N denotes the group of characters of G , then we have $\mathbf{K}_0(\mathcal{X}) \simeq \mathbf{K}_0(X) \otimes_{\mathbf{Z}} \mathbf{Z}[N]$ (see [SGA3, Exp. I, 4.7.3], as in [Tho1, Lem 4.14]). Thus we may write $\alpha = \sum_i a_i \otimes b_i$, where $a_i \in \mathbf{K}_0(X)$ and b_i are pairwise distinct elements of N . Restricting α along $BG_{k(x)} \rightarrow \mathcal{X}$ and identifying $\mathbf{K}_0(BG_{k(x)}) \simeq \mathbf{Z}[N]$, we get $\sum_i \mathrm{rk}(a_i) \cdot b_i = 0$ in $\mathbf{Z}[N]$, and hence $\mathrm{rk}(a_i) = 0$ for all i . Thus by [SGA6, Exp. VI, Prop. 6.1], a_i are all nilpotent as elements of $\mathbf{K}_0(X)$. In particular, α is also nilpotent as claimed.

¹³One could skip the reductions above by simply observing that the argument of *loc. cit.* goes through without the locally separatedness assumptions.

3.5. The equivariant master theorem. We formulate a G -equivariant version of condition **(L)**:

Definition 3.13. Let G be an fppf group algebraic space over k acting on $X \in \text{Stk}_k$, and $\Sigma \subseteq \text{Pic}(BG)$ a subset. We introduce the following condition on Σ :

(L_G) For every geometric point of X there exists a rank one G -representation $\mathcal{L}(x) \in \Sigma$ such that the $\text{St}_X^G(x)$ -representation $\mathcal{L}(x)|_{\text{BSt}_X^G(x)}$ is trivial.

Here $\text{St}_X^G(x)$ is the G -stabilizer at the point x (see Definition 1.4).

Specializing Theorem 3.5 to the case of quotient stacks yields the following equivariant version of the master theorem. As in Notation 3.1, $(-)\Sigma\text{-loc}$ denotes $(-)\langle * \rangle [c_1(\Sigma^{-1})]$.

Corollary 3.14. *Let $S \in \text{Stk}_k$ and G an fppf group algebraic space over k acting on a 1-Artin $X \in \text{Stk}_S$ with affine stabilizers. Let $\Sigma \subseteq \text{Pic}(BG)$ be a subset which satisfies condition **(L_G)** for X . Then we have*

$$\mathbf{C}_{\bullet}^{\text{BM}}([X/G]_{/S})_{\Sigma\text{-loc}} = 0$$

and in particular $\mathbf{C}_{\bullet}^{\text{BM},G}(X)_{\Sigma\text{-loc}} = 0$.

Proof. Apply Theorem 3.5 to the quotient stack $\mathcal{X} = [X/G]$ (with $\Sigma \subseteq \text{Pic}(\mathcal{X})$ the inverse image of Σ along $\mathcal{X} = [X/G] \rightarrow BG$). Since there is a commutative square

$$\begin{array}{ccc} \text{BAut}_{\mathcal{X}}(x) & \longrightarrow & \text{BSt}_X^G(x) \\ \downarrow & & \downarrow \\ \mathcal{X} & \longrightarrow & BG, \end{array}$$

triviality of a line bundle over $\text{BSt}_X^G(x)$ implies triviality over $\text{BAut}_{\mathcal{X}}(x)$. \square

Corollary 3.15. *Let G be an fppf group algebraic space over k . Let $S \in \text{Stk}_k$ and $i : Z \rightarrow X$ a G -equivariant closed immersion over S where $Z, X \in \text{Stk}_S$ are quasi-compact 1-Artin with affine stabilizers. Let $\Sigma \subseteq \text{Pic}(BG)$ be a subset which satisfies condition **(L_G)** for $X \setminus Z$. Then the direct image map*

$$i_* : \mathbf{C}_{\bullet}^{\text{BM}}([Z/G]_{/S})_{\Sigma\text{-loc}} \rightarrow \mathbf{C}_{\bullet}^{\text{BM}}([X/G]_{/S})_{\Sigma\text{-loc}}$$

is invertible.

We similarly have an equivariant version of conditions **(K)**:

Variante 3.16. Let G act on $X \in \text{Stk}_k$ and $\Sigma \subseteq \text{R}(G) \simeq \text{K}_0(BG)$ a subset.

(K_G) For every geometric point x of X we have

$$\text{K}_0(\text{BSt}_X^G(x))[\Sigma^{-1}] = 0.$$

For Λ rational, we may then specialize Theorem 3.10 to the equivariant case as above.

3.6. Base change.

Lemma 3.17. *Let $\mathcal{X}, \mathcal{X}' \in \text{Stk}_k$ be 1-Artin and let $f : \mathcal{X}' \rightarrow \mathcal{X}$ be a morphism. Let $\Sigma \subseteq \text{Pic}(\mathcal{X})$ be a subset and denote by $\Sigma' \subseteq \pi_0 \mathbf{C}^\bullet(\mathcal{X}') \langle * \rangle$ its image by f^* . If Σ satisfies condition (L) for \mathcal{X} , then Σ' satisfies condition (L) for \mathcal{X}' .*

Proof. Let x' be a geometric point of \mathcal{X}' and consider its image $x = f(x')$ in \mathcal{X} . By assumption, there exists an invertible sheaf $\mathcal{L}(x) \in \Sigma$ with $c_1(\mathcal{L}(x))|_{B\text{Aut}_{\mathcal{X}}(x)} = 0$. Then its inverse image $\mathcal{L}(x') := f^* \mathcal{L}(x)$ belongs to Σ' . Since there is a morphism of group schemes $\text{Aut}_{\mathcal{X}'}(x') \rightarrow \text{Aut}_{\mathcal{X}}(x)$, we also have $c_1(\mathcal{L}(x'))|_{B\text{Aut}_{\mathcal{X}'}(x')} = 0$. \square

Corollary 3.18. *Let G be an fppf group algebraic space over k acting on $X, X' \in \text{Stk}_k$ which are 1-Artin and let $f : X' \rightarrow X$ be a G -equivariant morphism. If $\Sigma \subseteq \text{Pic}(BG)$ satisfies condition (L_G) for X , then it also satisfies condition (L_G) for X' .*

Combining this with equivariant concentration (Theorem 3.15) yields:

Corollary 3.19. *Let G be an fppf group algebraic space over k . Let $S \in \text{Stk}_k$, $Z, X \in \text{Stk}_S$ quasi-compact 1-Artin with affine stabilizers, and $i : Z \rightarrow X$ a G -equivariant closed immersion over S . Let $\Sigma \subseteq \text{Pic}(BG)$ be a subset which satisfies condition (L_G) for $X \setminus Z$. Then for every morphism $f : X' \rightarrow X$, direct image along the base change $i' : Z' = Z \times_X X' \rightarrow X'$ induces an isomorphism*

$$i'_* : \mathbf{C}_{\bullet}^{\text{BM},G}(Z')_{\Sigma\text{-loc}} \rightarrow \mathbf{C}_{\bullet}^{\text{BM},G}(X')_{\Sigma\text{-loc}}.$$

Construction 3.20. Let G be an fppf group algebraic space over k with connected fibres, acting on a finite type 1-Artin $X \in \text{Stk}_k$ with affine stabilizers. Suppose we are given a G -equivariant morphism $\pi : X \rightarrow M$ to a finite type algebraic space $M \in \text{Stk}_k$. Denote by M^G the fixed locus of M (see Theorem 1.12(i)), and form the cartesian square

$$\begin{array}{ccc} M^G \times_M X & \longrightarrow & X \\ \downarrow & & \downarrow \pi \\ M^G & \longrightarrow & M. \end{array}$$

Corollary 3.21. *Let the notation be as in Construction 3.20. For any subset $\Sigma \subseteq \text{Pic}(BG)$, condition (L_G) for $M \setminus M^G$ implies condition (L_G) for $X \setminus (M^G \times_M X)$. In particular, the direct image map*

$$i_* : \mathbf{C}_{\bullet}^{\text{BM},G}(M^G \times_M X)_{\Sigma\text{-loc}} \rightarrow \mathbf{C}_{\bullet}^{\text{BM},G}(X)_{\Sigma\text{-loc}}$$

is invertible.

Remark 3.22. For example, if $G = T$ is diagonalizable then we may take Σ to be the set of nontrivial rank one representations $\mathcal{L} \in \text{Pic}(BT)$. Compare [Jos, Prop. 6.9] for G -theory of smooth Deligne–Mumford stacks with torus action over an algebraically closed field.

Remark 3.23. For example, if X has finite inertia (e.g. it is separated and Deligne–Mumford) then it admits a coarse moduli space M that is of

finite type [KM]. Moreover, the G -action automatically descends to M by universal properties, in such a way that $\pi : X \rightarrow M$ is equivariant. Similarly, if X admits a good or adequate moduli space $\pi : X \rightarrow M$ in the sense of Alper [Alp1, Alp2] and is of finite type, then M is of finite type [Alp2, Thm. 6.3.3].

3.7. Non-quasi-compact stacks. Let \mathcal{X} be a 1-Artin stack which has affine stabilizers but is only *locally* of finite type over k , and let $\Sigma \subseteq \text{Pic}(\mathcal{X})$. Unfortunately, concentration in the form of Theorem 3.5 does not hold for such \mathcal{X} : we may have

$$C_{\bullet}^{\text{BM}}(\mathcal{X})_{\text{loc}} \neq 0$$

even when Σ satisfies condition (L) for \mathcal{X} . In fact, we have simple counterexamples to Theorem 4.1 even for torus actions (and there are only finitely many T -stabilizer groups appearing).

Example 3.24. For every integer $n \geq 0$, consider the weight 1 scaling action of $T = \mathbf{G}_m$ on $U_n = \mathbf{A}^n \setminus 0$ and the element

$$\alpha_n = t \cap [U_n] \in C_{\bullet}^{\text{BM},T}(U_n)\langle -n+1 \rangle \simeq C_{\bullet}^{\text{BM}}(\mathbf{P}_k^{n-1})\langle -n+1 \rangle$$

where

$$t \in C_T^{\bullet}(\text{Spec}(k))\langle 1 \rangle \simeq C^{\bullet}(\mathbf{P}_k^{\infty})\langle 1 \rangle$$

is the first Chern class of the tautological line bundle on BT , and the isomorphism is [KR1, Cor. 6.1]. By the projective bundle formula, we have $t^i \cap \alpha_n = 0$ if and only if $i \geq n-1$. Thus if $X = \coprod_n U_n$, the element

$$\alpha = (\alpha_n)_n \in \widehat{C}_{\bullet}^{\text{BM},T}(X)\langle * \rangle \simeq \prod_{i \in \mathbf{Z}} C_{\bullet}^{\text{BM},T}(X)\langle i \rangle \simeq \prod_{i \in \mathbf{Z}} \prod_{n \geq 1} C_{\bullet}^{\text{BM},T}(U_n)\langle i \rangle$$

does not vanish after inverting t . Similarly, let $Y = \coprod_n U_n \times W_n$ where W_n is smooth 1-Artin of finite type over k , of pure dimension $-n$ with trivial T -action. Then $\beta_n = t \cap [U_n \times W_n] \in C_{\bullet}^{\text{BM},T}(U_n \times W_n)\langle 1 \rangle$, and the element

$$\beta = (\beta_n)_n \in C_{\bullet}^{\text{BM},T}(Y)\langle 1 \rangle \simeq \prod_n C_{\bullet}^{\text{BM},T}(U_n \times W_n)\langle 1 \rangle$$

does not vanish after inverting t .

In this subsection we give a definition of “localized Borel–Moore homology”

$$C_{\bullet}^{\text{BM}}(\mathcal{X})_{\text{loc}}$$

which is the same as the localization of Borel–Moore homology when \mathcal{X} is quasi-compact, but for which $C_{\bullet}^{\text{BM}}(\mathcal{X})_{\text{loc}} = 0$ still holds under condition (L). If we write \mathcal{X} as a filtered union of quasi-compact opens $\{\mathcal{X}_{\alpha}\}_{\alpha}$, then we will see $C_{\bullet}^{\text{BM}}(\mathcal{X})_{\text{loc}} \simeq \varprojlim_{\alpha} C_{\bullet}^{\text{BM}}(\mathcal{X}_{\alpha})_{\text{loc}}$.

Definition 3.25. Let $S \in \text{Stk}_k$, $\mathcal{X} \in \text{Stk}_S$, and $\Sigma \subseteq \text{Pic}(\mathcal{X})$. We define

$$C_{\bullet}^{\text{BM}}(\mathcal{X}/S)_{\text{loc}} := C_{\bullet}^{\text{BM}}(\mathcal{X}/S)_{\Sigma\text{-loc}} := \varprojlim_{\mathcal{X}' \in \mathcal{X}} C_{\bullet}^{\text{BM}}(\mathcal{X}'/S)\langle * \rangle [c_1(\Sigma^{-1})]$$

where the homotopy limit is taken over quasi-compact opens $\mathcal{X}' \subseteq \mathcal{X}$ (with restriction maps as the transition arrows).

Since condition (L) is stable under restriction to opens, the following non-quasi-compact generalization of Theorem 3.5 is immediate from the definition:

Theorem 3.26. *Let $S \in \text{Stk}_k$, $\mathcal{X} \in \text{Stk}_S$ which is 1-Artin with affine stabilizers, and $\Sigma \subseteq \text{Pic}(\mathcal{X})$. If Σ satisfies condition (L) for \mathcal{X} , then we have*

$$C_{\bullet}^{\text{BM}}(\mathcal{X}/S)_{\text{loc}} = 0.$$

We next note that this definition is compatible with Notation 3.1.

Proposition 3.27. *Let $S \in \text{Stk}_k$, $\mathcal{X} \in \text{Stk}_S$, and $\Sigma \subseteq \text{Pic}(\mathcal{X})$. If \mathcal{X} is a filtered union of quasi-compact opens $\{\mathcal{X}_{\alpha}\}_{\alpha}$, then there is a canonical isomorphism*

$$C_{\bullet}^{\text{BM}}(\mathcal{X}/S)_{\text{loc}} \simeq \varprojlim_{\alpha} C_{\bullet}^{\text{BM}}(\mathcal{X}_{\alpha}/S)_{\text{loc}}.$$

Proof. Consider the following commutative square:

$$\begin{array}{ccc} \varprojlim_{\mathcal{X}'} C_{\bullet}^{\text{BM}}(\mathcal{X}')\langle *\rangle[\Sigma^{-1}] & \longrightarrow & \varprojlim_{\alpha} C_{\bullet}^{\text{BM}}(\mathcal{X}_{\alpha})\langle *\rangle[\Sigma^{-1}] \\ \downarrow & & \downarrow \\ \varprojlim_{\mathcal{X}'} \varprojlim_{\alpha} C_{\bullet}^{\text{BM}}(\mathcal{X}_{\alpha} \cap \mathcal{X}')\langle *\rangle[\Sigma^{-1}] & \xlongequal{\quad} & \varprojlim_{\alpha} \varprojlim_{\mathcal{X}'} C_{\bullet}^{\text{BM}}(\mathcal{X}_{\alpha} \cap \mathcal{X}')\langle *\rangle[\Sigma^{-1}] \end{array}$$

where we omit the “ $/S$ ” from the notation for simplicity. The upper horizontal arrow comes from the fact that each \mathcal{X}_{α} is quasi-compact. The vertical arrows are induced by restriction. The right-hand one is invertible because $\mathcal{X}_{\alpha} \hookrightarrow \mathcal{X}$ is cofinal in $\{\mathcal{X} \cap \mathcal{X}_{\alpha} \hookrightarrow \mathcal{X}\}_{\alpha}$. The left-hand one is invertible because \mathcal{X}' is quasi-compact: we may choose a finite subset of opens in $(\mathcal{X}_{\alpha} \cap \mathcal{X}')_{\alpha}$ which cover \mathcal{X}' , and then by filteredness there exists some large enough index β such that $\mathcal{X}_{\beta} \cap \mathcal{X}' = \mathcal{X}'$. \square

Corollary 3.28. *Let $S \in \text{Stk}_k$, $\mathcal{X} \in \text{Stk}_S$, and $\Sigma \subseteq \text{Pic}(\mathcal{X})$. If \mathcal{X} is quasi-compact then there is a canonical isomorphism*

$$C_{\bullet}^{\text{BM}}(\mathcal{X}/S)_{\text{loc}} \simeq C_{\bullet}^{\text{BM}}(\mathcal{X}/S)\langle *\rangle[c_1(\Sigma)^{-1}].$$

Note also that $C_{\bullet}^{\text{BM}}(-)_{\text{loc}}$ still satisfies the localization triangle:

Proposition 3.29. *Let $S \in \text{Stk}_k$, $\mathcal{X} \in \text{Stk}_S$, and $\Sigma \subseteq \text{Pic}(\mathcal{X})$. Then for every closed immersion $i : \mathcal{Z} \hookrightarrow \mathcal{X}$ with open complement $j : \mathcal{U} \hookrightarrow \mathcal{X}$, we have a canonical exact triangle*

$$C_{\bullet}^{\text{BM}}(\mathcal{Z}/S)_{\text{loc}} \xrightarrow{i_*} C_{\bullet}^{\text{BM}}(\mathcal{X}/S)_{\text{loc}} \xrightarrow{j^!} C_{\bullet}^{\text{BM}}(\mathcal{U}/S)_{\text{loc}}.$$

Proof. Write \mathcal{X} as a filtered union of quasi-compact opens $\{\mathcal{X}_{\alpha}\}_{\alpha}$ (e.g. take the partially ordered set of all quasi-compact opens). For each α we have the localization triangle

$$C_{\bullet}^{\text{BM}}(\mathcal{Z} \cap \mathcal{X}_{\alpha}/S)\langle *\rangle[\Sigma^{-1}] \rightarrow C_{\bullet}^{\text{BM}}(\mathcal{X}_{\alpha}/S)\langle *\rangle[\Sigma^{-1}] \rightarrow C_{\bullet}^{\text{BM}}(\mathcal{U} \cap \mathcal{X}_{\alpha}/S)\langle *\rangle[\Sigma^{-1}]$$

since localization preserves exact triangles (as an exact functor). Passing to the homotopy limit (which is also exact) thus yields the claim by Corollary 3.27. \square

As before we may now specialize to the equivariant case. Let $S \in \text{Stk}_k$, G be an fppf group algebraic space over k which acts on a 1-Artin $X \in \text{Stk}_S$ with affine stabilizers, and $\Sigma \subseteq \text{Pic}(BG)$ a subset. We have

$$C_{\bullet}^{\text{BM}}([X/G]_{/S})_{\text{loc}} = \varprojlim_{X' \subseteq X} C_{\bullet}^{\text{BM}}([X'/G]_{/S}) \langle * \rangle [c_1(\Sigma)^{-1}]$$

where the limit is taken over quasi-compact G -invariant opens $X' \subseteq X$. We get the following non-quasi-compact version of Corollary 3.14:

Corollary 3.30. *If $\Sigma \subseteq \text{Pic}(BG)$ satisfies condition (L_G) for X , then we have*

$$C_{\bullet}^{\text{BM}}([X/G]_{/S})_{\Sigma\text{-loc}} = 0.$$

For \mathbf{D} rational, we have a parallel version of the above story for any $\Sigma \subseteq K_0(\mathcal{X})$, so that we get a non-quasi-compact version of Theorem 3.10 under condition (L) .

4. TORUS CONCENTRATION

In this section we fix a split torus T over k , and we assume that k is noetherian with no nontrivial idempotents.

4.1. The statement.

Theorem 4.1. *Let T act on a 1-Artin $X \in \text{Stk}_k$. Let $Z \subseteq X$ be a T -invariant closed substack such that for every point x of the complement $X \setminus Z$, $\text{St}_X^T(x)$ is properly contained in $T_{k(x)}$. Then we have:*

- (i) *Direct image along the closed immersion $i : Z \rightarrow X$ induces an isomorphism*

$$i_* : C_{\bullet}^{\text{BM},T}(Z)_{\Sigma\text{-loc}} \rightarrow C_{\bullet}^{\text{BM},T}(X)_{\Sigma\text{-loc}}$$

of $C^{\bullet}(BT)_{\Sigma\text{-loc}}$ -modules, where $\Sigma \subseteq \text{Pic}(BT)$ is the set of nontrivial line bundles on BT .

- (ii) *If X is quasi-compact then there exists a finite subset $\Sigma_0 \subseteq \Sigma$ such that we may replace Σ by Σ_0 in claim (i).*

Remark 4.2. In Theorem 4.1 the condition that $\text{St}_X^T(x) \not\subseteq T_{k(x)}$ is equivalent to the condition that $\dim(\text{St}_X^T(x)) < \dim(T_{k(x)})$, since $T_{k(x)}$ is irreducible.

Remark 4.3. Given a subgroup $H \subsetneq T$, consider the (diagonalizable) quotient $K = T/H$. Choose a nontrivial character of K , corresponding to a one-dimensional representation of K . The T -representation obtained by restriction may be regarded as a line bundle \mathcal{L} on BT whose restriction $\mathcal{L}|_{BH}$ is trivial.

Example 4.4. Suppose that the fixed locus X^T (Definition 1.9) is empty, i.e., for every geometric point x of X the T -stabilizer $\text{St}_X^T(x)$ is not equal to $T_{k(x)}$. Consider the set \mathfrak{G} of subgroups $H \subseteq T$ for which there exists a geometric point x of X such that the T -stabilizer $\text{St}_X^T(x)$ is equal to the base

change $H_{k(x)}$. For every $H \in \mathfrak{G}$ let $\mathcal{L}(H)$ be a nontrivial line bundle on BT such that $\mathcal{L}(H)|_{BH}$ is trivial (as in Remark 4.3). Then $\Sigma_0 = \{\mathcal{L}(H)\}_{H \in \mathfrak{G}}$ satisfies condition (L_T) for X .

Proposition 4.5. *Let T act on a quasi-compact 1-Artin $X \in \text{Stk}_k$. Then there exists a nonempty T -invariant open of X whose T -stabilizers are constant. In particular, the set \mathfrak{G} (Example 4.4) is finite.*

Proof. Note first of all that the second statement follows from the first by noetherian induction.

For the main statement, we may as well assume that X is reduced (since the statement only involves geometric points). For X an algebraic space, the statement follows immediately from Proposition 3.11. In general we argue as follows. By [SP, Tag 06QJ] and generic flatness of the morphism $[I_X/T] \rightarrow [X/T]$ (quotient of the projection $I_X \rightarrow X$), there exists a nonempty T -invariant open $X_0 \subseteq X$ such that X_0 is a gerbe (with respect to a flat finitely presented group algebraic space) over an algebraic space. Note also that this algebraic space is locally of finite type over k (since X is locally a trivial gerbe over it), hence of finite type over k since it is quasi-compact (because X is). Replacing X by X_0 we may therefore assume that X is a gerbe over an algebraic space M of finite type over k .

The T -action on X descends along $\pi : X \rightarrow M$, for example because it is a coarse moduli space. By the algebraic space case, it will suffice to show that $\pi : X \rightarrow M$ is T -stabilizer-preserving, i.e., for every geometric point x of X the canonical morphism

$$\text{St}_X^T(x) \rightarrow \text{St}_M^T(\pi(x))$$

is invertible. Since this is a homomorphism of subgroups of $T_{k(x)}$, it is enough to show surjectivity. A geometric point of $\text{St}_M^T(\pi(x))$ is a geometric point t of T such that $t \cdot \pi(x) = \pi(x)$. But this means precisely that there is an identification $t \cdot x \simeq x$ as geometric points of X , since on geometric points π exhibits M as the set of connected components of X . \square

Proof of Theorem 4.1. Let $\Sigma_0 \subseteq \Sigma$ be as in Example 4.4; when X is quasi-compact this is finite by Proposition 4.5. Thus it will suffice to show that i_* becomes invertible after inverting $c_1(\Sigma_0)$. By construction, Σ_0 satisfies condition (L_T) for $X \setminus Z$, so the claim follows directly from Corollary 3.30 at least when X has affine stabilizers.

Otherwise, by the localization triangle we may replace X with $X \setminus Z$ and thereby reduce to showing that $C_{\bullet}^{\text{BM}, T}(X)_{\text{loc}} = 0$ when Σ_0 satisfies condition (L_T) for X . As in the proof of Proposition 4.5, using the localization triangle again, we may assume that there exists a T -equivariant stabilizer-preserving morphism $\pi : X \rightarrow M$ which exhibits X as a gerbe over an algebraic space M of finite type over k . Then Σ_0 satisfies condition (L_T) for M as it does for X and π is a surjection on geometric points. Since M is an algebraic space, it follows from Proposition 3.11 that there exists a nonempty T -invariant affine open U of M and an invertible sheaf $\mathcal{L} \in \Sigma_0$

such that $c_1(\mathcal{L})$ is nilpotent as an element of the ring $\pi_0 C_T^\bullet(U)\langle * \rangle$ (see proof of Theorem 3.5). Therefore $V = \pi^{-1}(U)$ is a nonempty T -invariant open in X such that $c_1(\mathcal{L})$ is nilpotent in $\pi_0 C_T^\bullet(V)\langle * \rangle$. In particular

$$C_{\bullet}^{\text{BM},T}(V)\langle * \rangle[c_1(\mathcal{L})^{-1}] = 0$$

as it is a module over $C_T^\bullet(V)\langle * \rangle[c_1(\mathcal{L})^{-1}] = 0$. The claim now follows by noetherian induction using the localization triangle. \square

4.2. Example: fixed loci. In [AKLPR, App. A] we defined a T -fixed locus $X^T \subseteq X$ for an Artin stack X . In terms of X^T , the condition in Theorem 4.1 is that the open complement $X \setminus Z$ is contained in $X \setminus X^T$. A priori, the substack $X^T \subseteq X$ may not be closed in general. Nevertheless we obtain the following variants of concentration using the reduced fixed locus (Definition 1.10) or homotopy fixed point stack (Definition 1.11) when X has finite stabilizers or is Deligne–Mumford, respectively.

Corollary 4.6. *Let the notation be as in Theorem 4.1 and suppose X has finite stabilizers. Denote by X_{red}^T the reduced T -fixed locus (Definition 1.10). Then direct image along the closed immersion $i : X_{\text{red}}^T \rightarrow X$ (Theorem 1.12(ii)) induces an isomorphism*

$$i_* : C_{\bullet}^{\text{BM},T}(X_{\text{red}}^T)_{\Sigma\text{-loc}} \rightarrow C_{\bullet}^{\text{BM},T}(X)_{\Sigma\text{-loc}}$$

of $C^\bullet(BT)_{\Sigma\text{-loc}}$ -modules.

In the Deligne–Mumford case we can alternatively use the homotopy fixed point stack.

Corollary 4.7. *Let the notation be as in Theorem 4.1 and suppose $X \in \text{Stk}_k$ is quasi-compact Deligne–Mumford. Choose a reparametrization $T' \twoheadrightarrow T$ as in Theorem 1.12(iii) and denote by $X^{hT'}$ the homotopy fixed point stack with respect to the induced T' -action (Definition 1.11). Then direct image along the closed immersion $i : X^{hT'} \rightarrow X$ (Theorem 1.12(iv)) induces an isomorphism*

$$i_* : C_{\bullet}^{\text{BM},T}(X^{hT'})_{\Sigma\text{-loc}} \rightarrow C_{\bullet}^{\text{BM},T}(X)_{\Sigma\text{-loc}}$$

of $C^\bullet(BT)_{\Sigma\text{-loc}}$ -modules.

4.3. Example: vector bundles and abelian cones. For future use we record the following example:

Proposition 4.8. *Let $X \in \text{Stk}_k$ be 1-Artin with finite stabilizers. Let $\mathcal{E} \in \text{Coh}^T(X) \simeq \text{Coh}(X \times BT)$ be a T -equivariant coherent sheaf on X . Write $E = \mathbf{V}_X(\mathcal{E})$ for the associated cone over X with T -action, and $E \setminus X$ for the complement of the zero section. If \mathcal{E} has no fixed part, i.e., $\mathcal{E}^{\text{fix}} \simeq 0$, then for every point v of E , we have $\text{St}_E^T(v) = T_{k(v)}$ if and only if v belongs to the zero section.*

Proof. Suppose first that X is the spectrum of a field. Since T acts trivially on X , the T -representation \mathcal{E} splits as a direct sum of 1-dimensional representations. Since \mathcal{E} has no fixed part, the latter have nonzero weights, so the claim is clear in this case.

Now consider the general case, i.e., X is 1-Artin with finite stabilizers. Let v be a field-valued point of E . It is clear that if v belongs to the image of the zero section then it has $\text{St}_E^T(v) = T_{k(v)}$. Conversely, suppose that $v \in E \setminus X$. To show that the inclusion $\text{St}_E^T(v) \subseteq T_{k(v)}$ is proper it will suffice to show that $\dim(\text{St}_E^T(v)) < \dim(T_{k(v)})$ (since the scheme $T_{k(v)}$ is irreducible). Note that E also has finite stabilizers, since it is affine over X . By the short exact sequence of group schemes over $k(v)$ (1.5)

$$1 \rightarrow \underline{\text{Aut}}_E(v) \rightarrow \underline{\text{Aut}}_{\mathcal{E}}(v) \rightarrow \text{St}_E^T(v) \rightarrow 1,$$

where $\mathcal{E} = [E/T]$ is the quotient, it will moreover suffice to show that $\dim(\underline{\text{Aut}}_{\mathcal{E}}(v)) < \dim(T_{k(v)})$.

Let x denote the projection of v in X , E_x the fibre of E over x , and $\mathcal{E}_x = [E_x/T]$ the quotient. We have another short exact sequence

$$1 \rightarrow \underline{\text{Aut}}_{\mathcal{E}_x}(v) \rightarrow \underline{\text{Aut}}_{\mathcal{E}}(v) \rightarrow \underline{\text{Aut}}_X(x),$$

by applying Remark 1.1 twice, to the representable morphism $\mathcal{E}_x \rightarrow \mathcal{E}$ and to the morphism $\mathcal{E} \rightarrow X \times BT \rightarrow X$. By the special case of our claim where the base X is $\text{Spec}(k(x))$, we have $\dim(\underline{\text{Aut}}_{\mathcal{E}_x}(v)) < \dim(T_{k(v)})$. Since $\underline{\text{Aut}}_X(x)$ is finite, it follows that $\dim(\underline{\text{Aut}}_{\mathcal{U}}(v)) < \dim(T_{k(v)})$ as claimed. \square

By Theorem 4.1 we get:

Corollary 4.9. *Let the notation be as in Proposition 4.8. Let $\Sigma \subseteq \text{Pic}(BT)$ be the subset of nontrivial line bundles. Then Σ satisfies condition (LT) for $E \setminus X$.*

Corollary 4.10. *Let $S \in \text{Stk}_k$, $X \in \text{Stk}_S$ with finite stabilizers. Let \mathcal{E} be a T -equivariant coherent sheaf on X with no fixed part, and write $E = \mathbf{V}_X(\mathcal{E})$. Then direct image along the zero section $0: X \rightarrow E$ induces an isomorphism of $\mathbf{C}^\bullet(BT)_{\Sigma\text{-loc}}$ -modules*

$$0_* : \mathbf{C}_{\bullet}^{\text{BM}}(X \times BT/S)_{\Sigma\text{-loc}} \rightarrow \mathbf{C}_{\bullet}^{\text{BM}}([E/T]/S)_{\Sigma\text{-loc}},$$

where Σ is as in Corollary 4.9.

Proof. Follows from Corollaries 4.9 and 3.15. \square

We warn the reader that the finite stabilizers assumption in Corollary 4.10 is necessary, as we can already see in the ‘‘simplest’’ example of an Artin stack with infinite stabilizers, i.e. $B\mathbf{G}_m$. Indeed, even the conclusion of Proposition 4.8 does not hold in this example:

Example 4.11. Let $Z = B\mathbf{G}_m$, $X = [\mathbf{A}^1/\mathbf{G}_m]$, and $i: Z \rightarrow X$ the closed immersion induced by $0: \text{Spec}(k) \rightarrow \mathbf{A}^1$. Let $T = \mathbf{G}_m$ act on X with weight 1, so that i is T -equivariant and induces a closed immersion

$$[Z/T] \simeq B\mathbf{G}_m \times BT \rightarrow [\mathbf{A}^1/\mathbf{G}_m \times T] \simeq [X/T].$$

Then the induced direct image map

$$\mathbf{C}_{\bullet}^{\text{BM},T}(B\mathbf{G}_m)_{\text{loc}} \rightarrow \mathbf{C}_{\bullet}^{\text{BM},T}([\mathbf{A}^1/\mathbf{G}_m])_{\text{loc}}$$

is *not* an isomorphism.

4.4. Example: Higgs sheaves on curves. Let k be a field and C a smooth proper and geometrically connected curve over k . Consider the moduli stack $\underline{\text{Coh}}(C)$ of coherent sheaves on C . This is a smooth 1-Artin stack locally of finite type over k .

Denote by $\underline{\text{Higgs}}$ the moduli stack of Higgs sheaves on C , i.e., the cotangent bundle of $\underline{\text{Coh}}(C)$ (which for us means the total space of the cotangent complex). This admits a canonical scaling action by the torus $T = \mathbf{G}_m$. Let $\mathbf{\Lambda}$ denote the closed substack parametrizing *nilpotent* Higgs sheaves, i.e., pairs (\mathcal{F}, θ) where $\mathcal{F} \in \text{Coh}$ and $\theta : \mathcal{F} \rightarrow \mathcal{F} \otimes K_C$ is nilpotent (with K_C the canonical bundle of C).

Theorem 4.12. *Let $\Sigma \subseteq \text{Pic}(BT)$ denote the set of nontrivial invertible sheaves on BT . Then the direct image map*

$$C_{\bullet}^{\text{BM},T}(\mathbf{\Lambda})_{\text{loc}} \rightarrow C_{\bullet}^{\text{BM},T}(\underline{\text{Higgs}})_{\text{loc}}$$

is invertible (where the localization is as in Subsect. 3.7).

Proof. Let \mathcal{F} be a Higgs sheaf over an algebraically closed extension field κ of k . By Theorem 4.1 it is enough to show that, if \mathcal{F} is not nilpotent, then the T -stabilizer at the corresponding geometric point of $\underline{\text{Higgs}}$ is a proper subgroup of T_{κ} . We are grateful to A. Minets for providing the following argument.

First note that if \mathcal{F} is not nilpotent, then either its maximal torsion Higgs subsheaf \mathcal{F}_0 is nilpotent or the quotient $\mathcal{F}' = \mathcal{F}/\mathcal{F}_0$ is nilpotent. Thus we may assume that \mathcal{F} is either torsion or locally free.

Recall that the moduli stack of locally free Higgs sheaves (= Higgs bundles) of rank r admits a canonical T -equivariant map (the Hitchin fibration) to the scheme $A_r = \bigoplus_{i=1}^r H^0(C, K_C^{\otimes i})$, where T acts on $H^0(C, K_C^{\otimes i})$ with weight i , by sending \mathcal{F} to the coefficients of the characteristic polynomial of its Higgs field. Moreover, a locally free Higgs sheaf \mathcal{F} is nilpotent if and only if its image in the Hitchin base is trivial. By Corollary 3.21, this implies the claim in the locally free case.

In the torsion case we argue as follows. Under the ‘‘BNR correspondence’’ (see [BNR] or [Sim, Lem. 6.8]), Higgs torsion sheaves \mathcal{F} are push-forwards of coherent sheaves on T_C^* of finite length. In particular, there is a ‘‘Hilbert–Chow’’ map from the moduli of torsion Higgs sheaves to $\text{Sym}^d(T_C^*)$, sending \mathcal{F} to the support (counted with multiplicities) of the corresponding sheaf on T_C^* . Now \mathcal{F} is nilpotent if and only if the corresponding sheaf on T_C^* is supported in the zero section, hence if and only if its image in $\text{Sym}^d(T_C^*)$ is fixed under the \mathbf{G}_m -scaling action. We conclude by Corollary 3.19. \square

Remark 4.13. For the substack of torsion Higgs sheaves, Theorem 4.12 recovers [Min, Cor. 4.3] by taking π_0 . In other words, we can regard the

result as a generalization of *loc. cit.* to arbitrary rank Higgs sheaves¹⁴ and to “higher” oriented Borel–Moore homology theories.

Remark 4.14. Theorem 4.12 admits an analogous statement for Higgs G -bundles, for a connected reductive group G , using the same Hitchin fibration argument. Similarly, there is a parabolic variant using the parabolic Hitchin fibration [Yok].

REFERENCES

- [AB] M. F. Atiyah, R. Bott, *The moment map and equivariant cohomology*. Topology **23** (1984), no. 1, 1–28.
- [AKLPR] D. Aranha, A. A. Khan, A. Latyntsev, H. Park, C. Ravi, *Virtual localization revisited*. [arXiv:2207.01652](https://arxiv.org/abs/2207.01652) (2022).
- [Alp1] J. Alper, *Good moduli spaces for Artin stacks*. Ann. Inst. Fourier **63** (2013), no. 6, 2349–2402.
- [Alp2] J. Alper, *Adequate moduli spaces and geometrically reductive group schemes*. Algebr. Geom. **1** (2014), no. 4, 489–531.
- [BNR] A. Beauville, M. S. Narasimhan, S. Ramanan, *Spectral curves and the generalized theta divisor*. J. Reine Angew. Math. **398** (1989), 169–179.
- [BP] Y. Bae, H. Park, *A comparison theorem for cycle theories for algebraic stacks*. In preparation.
- [Bot] R. Bott, *Vector fields and characteristic numbers*. Michigan Math. J. **14** (1967), 231–244.
- [CD] D.-C. Cisinski, F. Déglise, *Triangulated categories of mixed motives*. Springer Monogr. Math., to appear. [arXiv:0912.2110](https://arxiv.org/abs/0912.2110) (2012).
- [Dég] F. Déglise, *Orientation theory in arithmetic geometry*. In: K-Theory—Proceedings of the International Colloquium (Mumbai, 2016), 239–347, Hindustan Book Agency (2018).
- [EG1] D. Edidin, W. Graham, *Equivariant intersection theory*. Invent. Math. **131** (1998), no. 3, 595–644.
- [EG2] D. Edidin, W. Graham, *Localization in equivariant intersection theory and the Bott residue formula*. Am. J. Math. **120** (1998), no. 3, 619–636.
- [FSS] R. Fedorov, A. Soibelman, Y. Soibelman, *Motivic classes of moduli of Higgs bundles and moduli of bundles with connections*. [arXiv:1705.04890](https://arxiv.org/abs/1705.04890) (2017).
- [Ful] W. Fulton, *Intersection theory*. Springer (1998).
- [GP] T. Graber, R. Pandharipande, *Localization of virtual classes*. Invent. Math. **135** (1999), no. 2, 487–518.
- [Gai] D. Gaitsgory, *Notes on geometric Langlands: stacks*. Unpublished preprint, available at <https://people.math.harvard.edu/~gaitsgde/GL/Stackstext.pdf> (2011).
- [HML] J. Heller, J. Malagón-López, *Equivariant algebraic cobordism*. J. Reine Angew. Math. **684**, 87–112 (2013).
- [HR] J. Hall, D. Rydh, *Algebraic groups and compact generation of their derived categories of representations*. Indiana Univ. Math. J. **64**, no. 6, 1903–1923 (2015).
- [Jos] R. Joshua, *Riemann–Roch for algebraic stacks. I*. Compositio Math. **136** (2003), no. 2, 117–169.
- [KK] A. A. Khan, T. Kinjo, *3d cohomological Hall algebras for local surfaces*. Available at: <https://www.preschema.com/papers/dimredcoha.pdf> (2023).
- [KM] S. Keel, S. Mori, *Quotients by groupoids*. Ann. of Math. **145** (1997), no. 1, 193–213.

¹⁴A similar claim is made in [SS, Prop. 3.7], but the proof is not correct because the localization does not commute with cofiltered limits. This is related to the reason why the discussion in Subsect. 3.7 is necessary.

- [KP] Y.-H. Kiem, H. Park, *Virtual intersection theories*. Adv. Math. **388** (2021), Article ID 107858.
- [KPR] A. A. Khan, H. Park, C. Ravi, *Concentration and localization formulas in K-theory*. In preparation.
- [KR1] A. A. Khan, C. Ravi, *Equivariant generalized cohomology via stacks*. [arXiv:2209.07801](https://arxiv.org/abs/2209.07801) (2022).
- [KR2] A. A. Khan, C. Ravi, *Cohomological and categorical concentration theorems*. [arXiv:2308.01652](https://arxiv.org/abs/2308.01652) (2023).
- [Kha1] A. A. Khan, *Virtual fundamental classes for derived stacks I*. [arXiv:1909.01332](https://arxiv.org/abs/1909.01332) (2019).
- [Kha2] A. A. Khan, *Voevodsky’s criterion for constructible categories of coefficients*. Available at: <https://www.preschema.com/papers/six.pdf> (2021).
- [Kha3] A. A. Khan, *K-theory and G-theory of derived algebraic stacks*. Jpn. J. Math. **17** (2022), 1–61.
- [Kha4] A. A. Khan, *Weaves*. Available at: <https://www.preschema.com/papers/weaves.pdf> (2023).
- [Kre] A. Kresch, *Cycle groups for Artin stacks*. Invent. Math. **138** (1999), no. 3, 495–536.
- [Kri] A. Krishna, *Equivariant Cobordism for Torus Actions*. Adv. Math. **231** (2012), no. 5, 2858–2891.
- [LZ1] Y. Liu, W. Zheng, *Enhanced six operations and base change theorem for higher Artin stacks*. [arXiv:1211.5948](https://arxiv.org/abs/1211.5948) (2012).
- [LZ2] Y. Liu, W. Zheng, *Enhanced adic formalism and perverse t-structures for higher Artin stacks*. [arXiv:1404.1128](https://arxiv.org/abs/1404.1128) (2014).
- [Lur] J. Lurie, *Higher algebra*, version of 2017-09-18. <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [Min] A. Minets, *Cohomological Hall algebras for Higgs torsion sheaves, moduli of triples and sheaves on surfaces*. Sel. Math., New Ser. **26** (2020), no. 2, paper no. 30, 67 p.
- [SGA3] M. Demazure, A. Grothendieck (eds.), *Schémas en groupes* (SGA 3). Séminaire de Géométrie Algébrique du Bois-Marie 1962–1964. Lecture Notes in Mathematics **151**, **152**, **153**, Springer (1970).
- [SGA6] P. Berthelot, A. Grothendieck, L. Illusie, *Théorie des intersections and théorème de Riemann–Roch*. Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6). Lecture Notes in Mathematics **225**, Springer (1971).
- [SP] The Stacks Project. <https://stacks.math.columbia.edu>.
- [SS] F. Sala, O. Schiffmann, *Cohomological Hall algebra of Higgs sheaves on a curve*. Algebr. Geom. **7** (2020), no. 3, 346–376.
- [Sim] C. Simpson, *Moduli of representations of the fundamental group of a smooth projective variety II*. Publ. Math. IHÉS **80** (1994), 5–79.
- [Tho1] R. W. Thomason, *Comparison of equivariant algebraic and topological K-theory*. Duke Math. J. **53** (1986), no. 3, 795–825.
- [Tho2] R. W. Thomason, *Une formule de Lefschetz en K-théorie équivariante algébrique*. Duke Math. J. **68** (1992), no. 3, 447–462.
- [Toë] B. Toën, *Simplicial presheaves and derived algebraic geometry*, in: *Simplicial methods for operads and algebraic geometry*, 119–186, Adv. Courses Math. CRM Barcelona (2010).
- [Voe] V. Voevodsky, *\mathbf{A}^1 -homotopy theory*. Proceedings of the International Congress of Mathematicians, Vol. I (Berlin, 1998). Doc. Math. **1998**, Extra Vol. I, 579–604.
- [Yok] K. Yokogawa, *Compactification of moduli of parabolic sheaves and moduli of parabolic Higgs sheaves*. J. Math. Kyoto Univ. **33** (1993), 451–504.

Fakultät Mathematik, Universität Duisburg-Essen, Essen, 45127, Germany

Institute of Mathematics, Academia Sinica, Taipei, 10617, Taiwan

Mathematical Institute, University of Oxford, Oxford, OX2 6GG, United Kingdom

Department of Mathematical Sciences, Seoul National University, Seoul, 08826, Korea

School of Mathematics, Tata Institute of Fundamental Research, Mumbai, 400005 India